

Lecture notes on topological insulators

Ming-Che Chang

Department of Physics, National Taiwan Normal University, Taipei, Taiwan

(Dated: October 29, 2020)

Contents

I. Review of Berry phase	1
A. Non-degenerate energy level	1
B. Degenerate energy levels	3
C. Magnetic resonance of nuclear quadrupole	5
D. Numerical computation of Berry phase	6
References	8

I. REVIEW OF BERRY PHASE

We first review the **Berry phase** (also known as the **geometric phase**) for a *non-degenerate* energy level (**Abelian** case). The Berry phase for *degenerate* energy levels (**non-Abelian** case) is discussed in the second part.

Note: For the rest of the lecture notes, a matrix is written in the San-serif font, $M = (M_{\alpha\beta})$.

A. Non-degenerate energy level

Consider a system that varies slowly with parameters $\lambda(t)$. One example is a spin in a slowly varying magnetic field $\mathbf{B}(t)$. At each instant t , the time-independent Schrödinger equation is

$$H_{\lambda}|n, \lambda\rangle = \varepsilon_{n\lambda}|n, \lambda\rangle. \quad (1.1)$$

The eigenstates $|n, \lambda\rangle$ will be called the **snapshot states**.

Suppose energy level $\varepsilon_{n\lambda}$ is gapped from other levels with a minimal separation Δ_0 during the course of evolution, and the characteristic frequency of the changing parameter $\Omega_0 \ll \Delta_0/\hbar$, then according to the **quantum adiabatic theorem**, an initial state $|n, \lambda(0)\rangle$ would stay at the same level n (Fig. 1). After time t , it evolves to the snapshot state at $\lambda(t)$, multiplied by a *dynamical* phase factor,

$$|n, \lambda_0\rangle \rightarrow |\Psi_{n\lambda}(t)\rangle = e^{-\frac{i}{\hbar} \int_0^t dt' \varepsilon_{n\lambda(t')}} |n, \lambda(t)\rangle. \quad (1.2)$$

In principle, we are allowed to assign a λ -dependent phase $\gamma_n(\lambda(t))$ to the snapshot state $|n, \lambda(t)\rangle$, since the new state still satisfies Eq. (1.1). Thus, in general,

$$|\Psi_{n\lambda}(t)\rangle = e^{i\gamma_n(t)} e^{-\frac{i}{\hbar} \int_0^t dt' \varepsilon_{n\lambda(t')}} |n, \lambda(t)\rangle. \quad (1.3)$$

This extra phase had been deemed as removable since the early days of quantum mechanics, until Berry showed the contrary in 1984 (Berry, 1984).




FIG. 1 If a Hamiltonian varies slowly with time, then an electron at state $|n, \lambda\rangle$ would stay at the same level.

The phase γ_n is constrained by the time-dependent Schrödinger equation,

$$H|\Psi_{n\lambda}(t)\rangle = i\hbar \frac{\partial}{\partial t} |\Psi_{n\lambda}(t)\rangle. \quad (1.4)$$

Substitute Eq. (1.3) into Eq. (1.4), one has

$$\frac{d\gamma_n(t)}{dt} = i\langle n, \lambda | \frac{\partial}{\partial t} |n, \lambda\rangle \quad (1.5)$$

$$\rightarrow \gamma_n(t) = i \int_0^t dt' \langle n, \lambda | \frac{\partial}{\partial t'} |n, \lambda\rangle. \quad (1.6)$$

For a cyclic change of the parameters,

$$\lambda(0) \rightarrow \lambda(T) = \lambda(0), \quad (1.7)$$

one has,

$$\gamma_n(T) = i \oint_C d\lambda \cdot \langle n, \lambda | \frac{\partial}{\partial \lambda} |n, \lambda\rangle, \quad (1.8)$$

in which C is the loop traversed in the parameter space of λ . $\gamma_n(T)$, or $\gamma_n(C)$, is the **Berry phase** acquired by $|\Psi_n\rangle$ after one cycle in energy level n . It depends only on the geometry of the path C , but not on the rate of $\dot{\lambda}$, as long as it is slow enough so that inter-level transition can be neglected.

We will call the integrand in Eq. (1.8) the **Berry connection**,

$$\mathbf{A}_n(\lambda) \equiv i\langle n, \lambda | \frac{\partial}{\partial \lambda} |n, \lambda\rangle. \quad (1.9)$$

If one re-assigns a λ -dependent phase to the snapshot state,

$$|n, \lambda\rangle \rightarrow |n, \lambda'\rangle = e^{i\chi(\lambda)} |n, \lambda\rangle, \quad (1.10)$$




FIG. 2 A magnetic field sweeps slowly around the loop C .

where $e^{i\chi(\lambda)}$ is a single-valued function, then

$$\mathbf{A}_n(\lambda) \rightarrow \mathbf{A}'_n(\lambda) = \mathbf{A}_n(\lambda) - \frac{\partial \chi}{\partial \lambda}. \quad (1.11)$$

This is similar to the **gauge transformation** of a vector potential in electromagnetism. Before Berry's discovery, people thought that by solving $\partial \chi(\lambda)/\partial \lambda = \mathbf{A}_n(\lambda)$, $\mathbf{A}'_n(\lambda)$ can be set as zero and removed (Schiff, 1972). However, this is possible only if the loop integral for $\gamma_n(C)$ is zero. If not, then since $\gamma_n(C)$ is gauge invariant,

$$\begin{aligned} \gamma_n(C) &= \oint_C d\lambda \cdot \mathbf{A}'_n(\lambda) = \oint_C d\lambda \cdot \mathbf{A}_n(\lambda) \\ &\quad - \underbrace{\chi(\lambda(T)) - \chi(\lambda(0))}_{=2\pi \times \text{integer}}, \end{aligned} \quad (1.12)$$

it's impossible to remove $\mathbf{A}_n(\lambda)$ by gauge transformation.

If the parameter space is three-dimensional, then one can apply the **Stokes theorem** to write the Berry phase as a surface integral over the area S enclosed by the loop C ,

$$\gamma_n(C) = \int_S d^2\mathbf{a} \cdot \nabla_{\lambda} \times \mathbf{A}_n \quad (1.13)$$

$$= \int_S d^2\mathbf{a} \cdot \mathbf{F}_n, \quad (1.14)$$

in which $\mathbf{F}_n \equiv \nabla_{\lambda} \times \mathbf{A}_n$ is called the **Berry curvature**. In higher dimension, Stokes theorem remains valid, but needs to be written in the language of **differential form** (see App. ??).

For a small loop \square ,

$$\gamma_n(\square) \simeq d^2\mathbf{a} \cdot \mathbf{F}_n, \quad (1.15)$$

where $d^2\mathbf{a} = d^2a\hat{\mathbf{n}}$. Hence,

$$\mathbf{F}_n \cdot \hat{\mathbf{n}} \simeq \frac{\gamma_n(\square)}{d^2a}. \quad (1.16)$$

This becomes an equality when $d^2a \rightarrow 0$. That is, the Berry curvature at λ equals the ratio between the Berry phase for an infinitesimal loop around that point and the area of the loop.

Example: Consider a spin-1/2 electron in a magnetic field $\mathbf{B} = B\hat{\mathbf{n}}$ (see Fig. 2), where $\hat{\mathbf{n}} = (\sin\theta \cos\phi, \sin\theta \sin\phi, \cos\theta)$. The Hamiltonian is

$$H = -\boldsymbol{\mu} \cdot \mathbf{B} = \mu_B \boldsymbol{\sigma} \cdot \mathbf{B}, \quad \mu_B = \frac{e\hbar}{2mc}. \quad (1.17)$$

The eigen-energies are $\varepsilon_{\pm} = \pm\mu_B B$, and the eigenstates are

$$|\hat{\mathbf{n}}, +\rangle = \begin{pmatrix} \cos \frac{\theta}{2} \\ e^{i\phi} \sin \frac{\theta}{2} \end{pmatrix}, \quad |\hat{\mathbf{n}}, -\rangle = \begin{pmatrix} -e^{-i\phi} \sin \frac{\theta}{2} \\ \cos \frac{\theta}{2} \end{pmatrix}. \quad (1.18)$$

Note that $e^{i\alpha\pm\phi}|\hat{\mathbf{n}}, \pm\rangle$ are also eigenstates (α_{\pm} are not angle-dependent), but they are not single-valued if α_{\pm} is not an integer. We emphasize that, *when calculating the Berry phase, the snapshot states in Eq. (1.8) need to be single-valued*. Even if one adopts only single-valued states, the choice of basis is still not unique. For example, $|\hat{\mathbf{n}}, \pm\rangle' = e^{\mp i\phi}|\hat{\mathbf{n}}, \pm\rangle$ are also single-valued. You can check that $|\hat{\mathbf{n}}, \pm\rangle$ have ϕ -ambiguity at $\theta = \pi$ (but not at $\theta = 0$), while $|\hat{\mathbf{n}}, \pm\rangle'$ have ϕ -ambiguity at $\theta = 0$ (but not at $\theta = \pi$).

The magnetic field \mathbf{B} plays the role of the slowly varying parameters λ . The Berry connection can be calculated from Eq. (1.9) with the gradient operator,

$$\frac{\partial}{\partial \mathbf{B}} = \frac{\partial}{\partial B} \hat{e}_r + \frac{1}{B} \frac{\partial}{\partial \theta} \hat{e}_{\theta} + \frac{1}{B \sin \theta} \frac{\partial}{\partial \phi} \hat{e}_{\phi}. \quad (1.19)$$

It is left as an exercise to show that

$$\mathbf{A}_+(\mathbf{B}) = i \langle \mathbf{B}, + | \frac{\partial}{\partial \mathbf{B}} | \mathbf{B}, + \rangle \quad (1.20)$$

$$= -\frac{1}{2B} \frac{1 - \cos \theta}{\sin \theta} \hat{e}_{\phi}. \quad (1.21)$$

Similarly,

$$\mathbf{A}_-(\mathbf{B}) = \frac{1}{2B} \frac{1 - \cos \theta}{\sin \theta} \hat{e}_{\phi}. \quad (1.22)$$

Both $\mathbf{A}_{\pm}(\mathbf{B})$ are singular along $\theta = \pi$. These vector potentials are the same as that of a **magnetic monopole**, and the line of singularity corresponds to the **Dirac string** there.

The Berry curvature is,

$$\mathbf{F}_{\pm}(\mathbf{B}) = \nabla_{\mathbf{B}} \times \mathbf{A}_{\pm}(\mathbf{B}) = \mp \frac{1}{2} \frac{\hat{\mathbf{B}}}{B^2}. \quad (1.23)$$

This is the same as the field of a magnetic monopole at the origin with a magnetic charge $\mp 1/2$.

The Berry phase can be calculated using either the line integral of \mathbf{A}_{\pm} , or the surface integral of \mathbf{F}_{\pm} . For a loop in Fig. 2, the Berry phase is found to be,

$$\gamma_{\pm}(C) = \mp \frac{1}{2} \Omega(C), \quad (1.24)$$

where $\Omega(C) = 2\pi(1 - \cos\theta)$ is the solid angle, as seen from the origin, extended by the loop C . If the loop is lying on the x - y -plane, then the Berry phase can only be $\mp\pi$. That is, the state changes sign after a cyclic evolution. The phase in Eq. (1.24) has been confirmed by passing neutrons through a tube with a helical magnetic field (Fig. 3).




FIG. 3 Berry phase of an electron passing through a helical magnetic field (Bitter and Dubbers, 1987).

Note that, given $|\hat{n}, \pm\rangle' = e^{\mp i\phi} |\hat{n}, \pm\rangle$, the Berry connections calculated from the two basis differ by a gradient,

$$\mathbf{A}'_{\pm}(\mathbf{B}) = \mathbf{A}_{\pm}(\mathbf{B}) \pm \frac{\partial\phi}{\partial\mathbf{B}} \quad (1.25)$$

$$= \mathbf{A}_{\pm}(\mathbf{B}) \pm \frac{1}{B \sin\theta} \hat{e}_{\phi}. \quad (1.26)$$

Therefore,

$$\mathbf{A}'_{\pm}(\mathbf{B}) = \pm \frac{1}{2B} \frac{1 + \cos\theta}{\sin\theta} \hat{e}_{\phi}. \quad (1.27)$$

Both $\mathbf{A}'_{\pm}(\mathbf{B})$ are singular along $\theta = 0$. They generate the same Berry curvatures as those in Eq. (1.23).

The integral of \mathbf{F}_{\pm} over a closed surface enclosing the origin is quantized,

$$\frac{1}{2\pi} \int_{S_B^2} d^2\mathbf{a} \cdot \mathbf{F}_{\pm}(\mathbf{B}) = \mp 1. \quad (1.28)$$

This integer remains fixed no matter how the surface S_B^2 is deformed, as long as it is not torn up. It is sometimes called the **Berry index** (or the **topological charge**) of the degenerate point.

Finally, the analogy between the gauge structure of the Berry phase and that of the classical electromagnetism is summarized in Table I.

B. Degenerate energy levels

One can extend the analysis above to degenerate energy levels (Wilczek and Zee, 1984). The wave function now has multiple components for a given energy level. As a result, the Berry connection and the Berry curvature become matrix-valued vectors (or vector-valued matrices). For simplicity, we consider an energy level $\varepsilon_{n\lambda}$ with

TABLE I Analogy between electromagnetism and anholonomy

Electromagnetism	Quantum anholonomy
vector potential $\mathbf{A}(\mathbf{r})$	Berry connection $\mathbf{A}(\boldsymbol{\lambda})$
magnetic field $\mathbf{B}(\mathbf{r})$	Berry curvature $\mathbf{F}(\boldsymbol{\lambda})$
magnetic monopole	degenerate point
magnetic charge	Berry index
magnetic flux $\Phi(C)$	Berry phase $\gamma(C)$

only two (globally) degenerate eigenstates, $|n, 1, \boldsymbol{\lambda}\rangle$ and $|n, 2, \boldsymbol{\lambda}\rangle$. Again $\boldsymbol{\lambda}$ are slowly varying parameters. After time t , the states evolve to (compare with Eq. (1.3))

$$|\Psi_{n,1}(t)\rangle = e^{-\frac{i}{\hbar} \int_0^t dt' \varepsilon_{n\lambda}(t')} \quad (1.29)$$

$$\times (|n, 1, \boldsymbol{\lambda}(t)\rangle \Gamma_{11}(t) + |n, 2, \boldsymbol{\lambda}(t)\rangle \Gamma_{21}(t)),$$

$$|\Psi_{n,2}(t)\rangle = e^{-\frac{i}{\hbar} \int_0^t dt' \varepsilon_{n\lambda}(t')} \quad (1.30)$$

$$\times (|n, 1, \boldsymbol{\lambda}(t)\rangle \Gamma_{12}(t) + |n, 2, \boldsymbol{\lambda}(t)\rangle \Gamma_{22}(t)).$$

Or, ($\alpha, \beta = 1, 2$)

$$|\Psi_{n\beta}(t)\rangle = e^{-\frac{i}{\hbar} \int_0^t dt' \varepsilon_{n\lambda}(t')} \sum_{\alpha} |n\alpha\boldsymbol{\lambda}(t)\rangle \Gamma_{\alpha\beta}(t). \quad (1.31)$$

We call the matrix $\Gamma_{\alpha\beta}$ a **Berry rotation matrix**.

Substitute the states into the Schrödinger equation,

$$H|\Psi_{n\beta}(t)\rangle = i\hbar \frac{\partial}{\partial t} |\Psi_{n\beta}(t)\rangle, \quad (1.32)$$

one will get,

$$\frac{d\Gamma_{\alpha\beta}}{dt} = - \sum_{\gamma} \langle n\alpha\boldsymbol{\lambda} | \frac{\partial}{\partial t} | n\gamma\boldsymbol{\lambda} \rangle \Gamma_{\gamma\beta} \quad (1.33)$$

$$= i \sum_{\gamma} \dot{\boldsymbol{\lambda}}(t) \cdot \mathbf{A}_{\alpha\gamma}^{(n)}(\boldsymbol{\lambda}) \Gamma_{\gamma\beta}, \quad (1.34)$$

where

$$\mathbf{A}_{\alpha\beta}^{(n)}(\boldsymbol{\lambda}) \equiv i \langle n\alpha\boldsymbol{\lambda} | \frac{\partial}{\partial \boldsymbol{\lambda}} | n\beta\boldsymbol{\lambda} \rangle. \quad (1.35)$$

The Berry connection now becomes a matrix-valued vector, $\vec{\mathbf{A}}^{(n)}$. To simplify the index, we focus only on one degenerate eigenenergy and suppress the index n from now on.

To solve for the $U(2)$ Berry rotation matrix $\Gamma(t)$, first consider an infinitesimal dt ,

$$\Gamma(t + dt) = \Gamma(t) + idt \dot{\boldsymbol{\lambda}}(t) \cdot \vec{\mathbf{A}}(t) \Gamma(t) \quad (1.36)$$

$$\simeq e^{idt \dot{\boldsymbol{\lambda}}(t) \cdot \vec{\mathbf{A}}(t)} \Gamma(t) \quad (1.37)$$

A full path can be divided into small steps, so that

$$\Gamma(t) = \dots e^{id\boldsymbol{\lambda} \cdot \vec{\mathbf{A}}(\boldsymbol{\lambda}_1)} e^{id\boldsymbol{\lambda} \cdot \vec{\mathbf{A}}(\boldsymbol{\lambda}_0)} \Gamma(0) \quad (1.38)$$

$$\equiv P e^{i \int_{\boldsymbol{\lambda}_0}^{\boldsymbol{\lambda}(t)} d\boldsymbol{\lambda} \cdot \vec{\mathbf{A}}(\boldsymbol{\lambda})}, \Gamma(0) = 1, \quad (1.39)$$

in which P is a **path-ordering operator**. Because of the path-ordering, $\Gamma(t)$ usually can only be computed numerically (see Sec. I.D).

Sometimes one can see non-Abelian Berry connections being calculated for energy levels that are not globally degenerate (or not degenerate at all). In such cases, the dynamical phase factor in Eq. (1.31) needs be changed to $e^{-\frac{i}{\hbar} \int_0^t dt' \varepsilon_{n\alpha\lambda}(t')}$, and the right hand side of Eq. (1.34) is multiplied by $e^{-\frac{i}{\hbar} \int_0^t dt' (\varepsilon_{n\gamma\lambda} - \varepsilon_{n\alpha\lambda})}$. The Berry connection and Berry curvature can still be defined in the same manner, but the Berry rotation among these energy levels would be mixed with dynamical phase factors.

We now consider a gauge transformation,

$$|\alpha\lambda\rangle' = \sum_{\gamma} |\gamma\lambda\rangle U_{\gamma\alpha}(\lambda). \quad (1.40)$$

Its dual is

$$\langle\alpha\lambda|' = \sum_{\gamma} U_{\alpha\gamma}^{\dagger}(\lambda) \langle\gamma\lambda|. \quad (1.41)$$

Therefore,

$$\mathbf{A}'_{\alpha\beta}(\lambda) = i \langle n\alpha\lambda|' \frac{\partial}{\partial \lambda} |n\beta\lambda\rangle' \quad (1.42)$$

$$= U_{\alpha\gamma}^{\dagger} \mathbf{A}_{\gamma\delta} U_{\delta\beta} + i U_{\alpha\gamma}^{\dagger} \frac{\partial}{\partial \lambda} U_{\gamma\beta}. \quad (1.43)$$

Or,

$$\mathbf{A}'_k = \mathbf{U}^{\dagger} \mathbf{A}_k \mathbf{U} + i \mathbf{U}^{\dagger} \frac{\partial}{\partial \lambda_k} \mathbf{U}. \quad (1.44)$$

The phase factor

$$e^{i d\lambda \cdot \vec{A}'} \simeq 1 + i d\lambda \cdot \vec{A}' \quad (1.45)$$

$$= 1 + i d\lambda \cdot \mathbf{U}^{\dagger} \vec{A} \mathbf{U} - d\lambda \cdot \mathbf{U}^{\dagger} \frac{\partial}{\partial \lambda} \mathbf{U} \quad (1.46)$$

$$\simeq \mathbf{U}^{\dagger} \left(1 + i d\lambda \cdot \vec{A} \right) \left(\mathbf{U} - d\lambda \cdot \frac{\partial}{\partial \lambda} \mathbf{U} \right) \quad (1.47)$$

$$\simeq \mathbf{U}^{\dagger}(\lambda) e^{i d\lambda \cdot \vec{A}} \mathbf{U}(\lambda - d\lambda). \quad (1.48)$$

Therefore, after a closed path,

$$\Gamma'[C] = \mathbf{U}^{\dagger}(\lambda_0) \Gamma[C] \mathbf{U}(\lambda_0). \quad (1.49)$$

That is, the non-Abelian Berry rotation is **gauge covariant**.

Berry curvature \vec{F} is defined as the Berry rotation Γ_{\square} around an infinitesimal loop \square enclosing an area $d^2\mathbf{a}$, $\Gamma_{\square} = e^{i \vec{F}_{\square} \cdot d^2\mathbf{a}}$, or

$$i \vec{F}_{\square} \cdot \hat{\mathbf{n}} \equiv \frac{\Gamma_{\square} - 1}{d^2 a}, \quad (1.50)$$

where $\hat{\mathbf{n}}$ is the normal vector of the surface $d^2\mathbf{a}$.

The Berry rotation matrix is composed of 4 steps (see Fig. 4),

$$\begin{aligned} & \Gamma_{\square}(\lambda) \\ &= \Gamma(\lambda, \lambda + d\lambda_2) \Gamma(\lambda + d\lambda_2, \lambda + d\lambda_1 + d\lambda_2) \\ & \times \Gamma(\lambda + d\lambda_1 + d\lambda_2, \lambda + d\lambda_1) \Gamma(\lambda + d\lambda_1, \lambda). \end{aligned} \quad (1.51)$$




FIG. 4 A small loop in the parameter space of λ .

Using the approximation,

$$e^{\epsilon \mathbf{A}} e^{\epsilon \mathbf{B}} = e^{\epsilon(\mathbf{A}+\mathbf{B}) + \frac{\epsilon^2}{2} [\mathbf{A}, \mathbf{B}] + O(\epsilon^3)}, \quad (1.52)$$

one has

$$\begin{aligned} & \Gamma(\lambda + d\lambda_1 + d\lambda_2, \lambda + d\lambda_1) \Gamma(\lambda + d\lambda_1, \lambda) \\ &= e^{i \mathbf{A}_k(\lambda + d\lambda_1) d\lambda_{2k}} e^{i \mathbf{A}_\ell(\lambda) d\lambda_{1\ell}} \end{aligned} \quad (1.53)$$

$$\simeq e^{i(\mathbf{A}_k d\lambda_{2k} + \mathbf{A}_\ell d\lambda_{1\ell} + \partial_\ell \mathbf{A}_k d\lambda_{2k} d\lambda_{1\ell}) - \frac{1}{2} [\mathbf{A}_k, \mathbf{A}_\ell] d\lambda_{2k} d\lambda_{1\ell}}. \quad (1.54)$$

For the next two steps, it is convenient to adopt instead (see p. 239 of [Cheng and Li, 1988](#)),

$$\begin{aligned} & \Gamma^{-1}(\lambda + d\lambda_2, \lambda) \Gamma^{-1}(\lambda + d\lambda_1 + d\lambda_2, \lambda + d\lambda_2) \\ &= e^{-i \mathbf{A}_k(\lambda) d\lambda_{2k}} e^{-i \mathbf{A}_\ell(\lambda + d\lambda_2) d\lambda_{1\ell}} \end{aligned} \quad (1.55)$$

$$\simeq e^{-i(\mathbf{A}_k d\lambda_{2k} + \mathbf{A}_\ell d\lambda_{1\ell} + \partial_k \mathbf{A}_\ell d\lambda_{2k} d\lambda_{1\ell}) - \frac{1}{2} [\mathbf{A}_k, \mathbf{A}_\ell] d\lambda_{2k} d\lambda_{1\ell}} \quad (1.56)$$

Therefore,

$$\Gamma_{\square} = e^{-i(\partial_k \mathbf{A}_\ell - \partial_\ell \mathbf{A}_k - i[\mathbf{A}_k, \mathbf{A}_\ell]) d\lambda_{2k} d\lambda_{1\ell}} \quad (1.57)$$

$$= e^{i \mathbf{F}_{k\ell} d\lambda_{1k} d\lambda_{2\ell}}, \quad (1.58)$$

with the Berry curvature,

$$\mathbf{F}_{k\ell} = \partial_k \mathbf{A}_\ell - \partial_\ell \mathbf{A}_k - i[\mathbf{A}_k, \mathbf{A}_\ell]. \quad (1.59)$$

Note that $\mathbf{F}_{k\ell}$ is antisymmetric in k, ℓ , and

$$\mathbf{F}_{k\ell} d\lambda_{1k} d\lambda_{2\ell} = \frac{1}{2} \epsilon_{cab} \mathbf{F}_{ab} \epsilon_{ck\ell} d\lambda_{1k} d\lambda_{2\ell}, \quad (1.60)$$

in which $\frac{1}{2} \epsilon_{cab} \mathbf{F}_{ab} = \mathbf{F}_c$ is the vector equivalent of the antisymmetric matrix, and $\epsilon_{ck\ell} d\lambda_{1k} d\lambda_{2\ell} = (d^2\mathbf{a})_c$. Therefore,

$$\mathbf{F}_{k\ell} d\lambda_{1k} d\lambda_{2\ell} = \vec{F} \cdot d^2\mathbf{a}, \quad (1.61)$$

When written in the vectorial form, the Berry curvature is (cf. Eq. (1.59)),

$$\vec{F} = \nabla_{\lambda} \times \vec{A} - i \vec{A} \times \vec{A}. \quad (1.62)$$

Finally, under a gauge transformation,

$$\mathbf{F}_{k\ell} \rightarrow \mathbf{F}'_{k\ell} = \partial_k \mathbf{A}'_{\ell} - \partial_\ell \mathbf{A}'_k - i[\mathbf{A}'_k, \mathbf{A}'_{\ell}] \quad (1.63)$$

$$= \mathbf{U}^{\dagger} \mathbf{F}_{k\ell} \mathbf{U}. \quad (1.64)$$

That is, the non-Abelian Berry curvature is gauge covariant (similarly for \vec{F}).

C. Magnetic resonance of nuclear quadrupole

The interaction between a nuclear quadrupole with a magnetic field \mathbf{B} is (Zee, 1988),

$$\mathbf{H} = (\vec{\mathbf{J}} \cdot \mathbf{B})^2, \quad (1.65)$$

where $\vec{\mathbf{J}}$ is the spin operator for spin- j . This coupling is invariant under time reversal. The eigen-energies and eigenstates are

$$\mathbf{H}|\hat{B}, \pm m\rangle = (mB)^2|\hat{B}, \pm m\rangle, \quad m = -j, \dots, j. \quad (1.66)$$

Note that $|\hat{B}, \pm m\rangle$ are degenerate with the same energy $(mB)^2$.

Since $\sigma_\alpha^2 = 1$ ($\alpha = x, y, z$) for the Pauli matrices, the Hamiltonian is non-trivial only if the spin quantum number $j \geq 1$. We are interested in half-integer spins, so the next non-trivial case is $j = 3/2$. In this case, the angular momentum matrices are

$$\mathbf{J}_x = \begin{pmatrix} 0 & \sqrt{3}/2 & 0 & 0 \\ \sqrt{3}/2 & 0 & 1 & 0 \\ 0 & 1 & 0 & \sqrt{3}/2 \\ 0 & 0 & \sqrt{3}/2 & 0 \end{pmatrix}, \quad \mathbf{J}_y = \begin{pmatrix} 0 & -i\sqrt{3}/2 & 0 & 0 \\ i\sqrt{3}/2 & 0 & -i & 0 \\ 0 & i & 0 & -i\sqrt{3}/2 \\ 0 & 0 & i\sqrt{3}/2 & 0 \end{pmatrix}, \quad \mathbf{J}_z = \begin{pmatrix} 3/2 & 0 & 0 & 0 \\ 0 & 1/2 & 0 & 0 \\ 0 & 0 & -1/2 & 0 \\ 0 & 0 & 0 & -3/2 \end{pmatrix}. \quad (1.67)$$

The spectrum of \mathbf{H} is composed of 2 two-fold degenerate energy levels. When \mathbf{B} is along the direction in Fig. 2, the eigenstates are ($m = -3/2, \dots, 3/2$)

$$|\hat{B}, m\rangle = \mathbf{U}_z(\phi)\mathbf{U}_y(\theta)|m\rangle \quad (1.68)$$

$$= e^{-iJ_z\phi}e^{-iJ_y\theta}|m\rangle. \quad (1.69)$$

The Berry connection is

$$\begin{aligned} \mathbf{A}_{mm'} &= i\langle \hat{B}, m | \frac{\partial}{\partial \mathbf{B}} | \hat{B}, m' \rangle \\ &= \frac{i}{B} \langle m | \mathbf{U}_y^\dagger(\theta)\mathbf{U}_z^\dagger(\phi) \left(\frac{\partial}{\partial \theta} \hat{e}_\theta + \frac{1}{\sin \theta} \frac{\partial}{\partial \phi} \hat{e}_\phi \right) \mathbf{U}_z(\phi)\mathbf{U}_y(\theta) | m' \rangle. \end{aligned} \quad (1.70)$$

With the help of the following equations,

$$e^{iJ_y\theta}e^{iJ_z\phi}i\frac{\partial}{\partial \theta}e^{-iJ_z\phi}e^{-iJ_y\theta} = \mathbf{J}_y, \quad (1.71)$$

$$e^{iJ_y\theta}e^{iJ_z\phi}i\frac{\partial}{\partial \phi}e^{-iJ_z\phi}e^{-iJ_y\theta} = -\sin \theta \mathbf{J}_x + \cos \theta \mathbf{J}_z, \quad (1.72)$$

one will get

$$\vec{\mathbf{A}} = \mathbf{A}_\theta \hat{e}_\theta + \mathbf{A}_\phi \hat{e}_\phi, \quad (1.73)$$

where

$$\mathbf{A}_\theta = \frac{1}{B} \mathbf{J}_y, \quad (1.74)$$

$$\mathbf{A}_\phi = \frac{1}{B \sin \theta} (-\sin \theta \mathbf{J}_x + \cos \theta \mathbf{J}_z). \quad (1.75)$$

Note that \mathbf{A}_ϕ blows up at $\theta = 0, \pi$.

In the calculation above, the eigenstates $|\hat{B}, m\rangle = \mathbf{U}_z(\phi)\mathbf{U}_y(\theta)|m\rangle$ are not single-valued if $\phi \rightarrow \phi + 2\pi$. One way to remedy this is to use (see ?),

$$|\hat{B}, m\rangle = \mathbf{U}_z(\phi)\mathbf{U}_y(\theta)\mathbf{U}_z(-\phi)|m\rangle \quad (1.76)$$

$$= e^{im\phi}\mathbf{U}_z(\phi)\mathbf{U}_y(\theta)|m\rangle. \quad (1.77)$$

Also, at $\theta = 0$, $|\hat{B}, m\rangle \rightarrow |m\rangle$ without the ϕ -ambiguity.

Using this new basis, one would get

$$\mathbf{A}_\theta = \frac{1}{B} \begin{pmatrix} 0 & -i\sqrt{3}/2e^{-i\phi} & 0 & 0 \\ i\sqrt{3}/2e^{i\phi} & 0 & -ie^{-i\phi} & 0 \\ 0 & ie^{i\phi} & 0 & -i\sqrt{3}/2e^{-i\phi} \\ 0 & 0 & i\sqrt{3}/2e^{i\phi} & 0 \end{pmatrix}, \quad (1.78)$$

and

$$\mathbf{A}_\phi = \frac{1}{B \sin \theta} \begin{pmatrix} \frac{3}{2}(\cos \theta - 1) & -\frac{\sqrt{3}}{2} \sin \theta e^{-i\phi} & 0 & 0 \\ -\frac{\sqrt{3}}{2} \sin \theta e^{i\phi} & \frac{1}{2}(\cos \theta - 1) & -\sin \theta e^{-i\phi} & 0 \\ 0 & -\sin \theta e^{i\phi} & -\frac{1}{2}(\cos \theta - 1) & -\frac{\sqrt{3}}{2} \sin \theta e^{-i\phi} \\ 0 & 0 & -\frac{\sqrt{3}}{2} \sin \theta e^{i\phi} & -\frac{3}{2}(\cos \theta - 1) \end{pmatrix}. \quad (1.79)$$




FIG. 5 Typical structure energy levels near the primary gap of semiconductors with direct energy gap.

Note that A_ϕ now is finite at $\theta = 0$ (but $\theta = \pi$ remains a singularity).

The Berry connection for the degenerate $|m| = 3/2$ levels are ($P_{3/2}$ is a projection operator)

$$\begin{aligned} \vec{A}^{(3/2)} &= P_{3/2} (A_\theta \hat{e}_\theta + A_\phi \hat{e}_\phi) P_{3/2} \quad (1.80) \\ &= \frac{1}{B \sin \theta} \begin{pmatrix} \frac{3}{2}(\cos \theta - 1) & 0 \\ 0 & -\frac{3}{2}(\cos \theta - 1) \end{pmatrix} \hat{e}_\phi, \quad (1.81) \end{aligned}$$

which is diagonal (Abelian connection). Similarly, the Berry connection for the degenerate $|m| = 1/2$ levels are

$$\begin{aligned} \vec{A}^{(1/2)} &= P_{1/2} (A_\theta \hat{e}_\theta + A_\phi \hat{e}_\phi) P_{1/2} \quad (1.82) \\ &= \frac{1}{B} \begin{pmatrix} 0 & -ie^{-i\phi} \\ ie^{i\phi} & 0 \end{pmatrix} \hat{e}_\theta \quad (1.83) \end{aligned}$$

$$+ \frac{1}{B \sin \theta} \begin{pmatrix} \frac{1}{2}(\cos \theta - 1) & -\sin \theta e^{-i\phi} \\ -\sin \theta e^{i\phi} & -\frac{1}{2}(\cos \theta - 1) \end{pmatrix} \hat{e}_\phi,$$

which is not diagonal (non-Abelian connection).

The only non-zero component of the Berry curvature $F_{k\ell}(k, \ell = r, \theta, \phi)$ is $F_{\theta\phi}$,

$$F_{\theta\phi}^{(3/2)} = \left(\nabla \times \vec{A}^{(3/2)} \right)_r - i[A_\theta^{(3/2)}, A_\phi^{(3/2)}] \quad (1.84)$$

$$= -\frac{3}{2} \frac{1}{B^2} \sigma_3. \quad (1.85)$$

It is left as an exercise for the readers to show this, and that

$$F_{\theta\phi}^{(1/2)} = \frac{3}{2} \frac{1}{B^2} \sigma_3. \quad (1.86)$$

These Berry curvatures are the same for the two types of basis in Eqs. (1.68), (1.76). Note that the Berry rotation is the (path-ordered) line integral of \vec{A} , but not the area integral of \vec{F} , since the Stokes theorem fails in the non-Abelian case. Because of the extra $\pm 3/2, \pm 1/2$ in the diagonal elements of A_ϕ in Eq. (1.79), for a circular path

with fixed θ , the Berry phases differ by a sign to those calculated from Eq. (1.75). Eq. (1.79) is to be trusted since it is based on single-valued snapshot states.

If you calculate the Berry curvature using Eqs. (1.78), (1.79) directly without projection, then the result would be zero. This can be explained as follows: The Berry curvature in Eq. (1.59) can be written as,

$$\begin{aligned} (F_{\alpha\beta})_{kl} &= i \langle \partial_k \alpha | \partial_l \beta \rangle - i \sum_\gamma \langle \partial_k \alpha | \gamma \rangle \langle \gamma | \partial_l \beta \rangle \\ &\quad - (k \leftrightarrow l). \quad (1.87) \end{aligned}$$

Without projecting the Berry connection to a subspace, γ runs through bases of the whole Hilbert space. Thus, $\sum_\gamma |\gamma\rangle \langle \gamma| = 1$ and F_{kl} vanishes.

Two remarks are in order: First, the Berry phase would result in splitting of the spectra of the nuclear quadrupole resonance. Such a splitting has been confirmed in experiment (Tycko, 1987). A recent attempt of detecting the non-Abelian Berry rotation can be found in Li *et al.*, 2016.

Second, the quadrupole coupling Hamiltonian is similar to the effective Hamiltonian of heavy holes ($m = \pm 3/2$) and light holes ($m = \pm 1/2$) near $\mathbf{k} = 0$ in a semiconductor (Fig. 5). Under the four-band approximation, and neglecting anisotropy, the four-band **Luttinger Hamiltonian** is (see, e.g., Winkler, 2003),

$$H = \frac{\hbar^2}{2m} \left[\left(\gamma_1 + \frac{5}{2} \gamma_2 \right) k^2 - 2\gamma_2 \left(\mathbf{k} \cdot \vec{J} \right)^2 \right], \quad (1.88)$$

where $\gamma_{1,2}$ are material-dependent parameters. The first term is multiplied by an identity matrix and has no effect on the Berry connection and curvature. The second term is essentially the same as the one in Eq. (1.65), if \mathbf{B} is replaced by \mathbf{k} . Therefore, this system has exactly the same Berry connection and Berry curvatures listed in Eqs. (1.78), (1.79), (1.85), and (1.86) (Murakami *et al.*, 2003).

D. Numerical computation of Berry phase

Numerical computation of Berry rotation using Eq. (1.39) can be implemented as follows: Divide a closed path C into N steps with points $\lambda_0, \lambda_1, \dots, \lambda_{N-1}$, and $\lambda_N = \lambda_0$ (Fig. 6). For each step, one has

$$\left[e^{i d\lambda \cdot \vec{A}(\lambda_\ell)} \right]_{\alpha\beta} \simeq \delta_{\alpha\beta} + i d\lambda \cdot \mathbf{A}_{\alpha\beta}(\lambda_\ell) \quad (1.89)$$

$$\begin{aligned} &= \langle \alpha \lambda_\ell | \beta \lambda_\ell \rangle - d\lambda \cdot \langle \alpha \lambda_\ell | \frac{\partial}{\partial \lambda} | \beta \lambda_\ell \rangle \\ &= \langle \alpha \lambda_\ell | \beta \lambda_\ell \rangle + d\lambda \cdot \langle \frac{\partial}{\partial \lambda} \alpha \lambda_\ell | \beta \lambda_\ell \rangle \\ &\simeq \langle \alpha \lambda_{\ell+1} | \beta \lambda_\ell \rangle. \quad (1.90) \end{aligned}$$

Thus (with the summation convention),

$$F_{\alpha\beta}(C) = \langle \alpha \lambda_0 | \beta_{N-1} \lambda_{N-1} \rangle \cdots \langle \beta_2 \lambda_2 | \beta_1 \lambda_1 \rangle \langle \beta_1 \lambda_1 | \beta \lambda_0 \rangle. \quad (1.91)$$




FIG. 6 Discretize a closed loop of path in parameter space.

When written in matrices, we have the Berry rotation,

$$\Gamma(C) = M_{N-1} \cdots M_1 M_0, \quad (1.92)$$

$$\text{where } (M_\ell)_{\alpha\beta} = \langle \alpha | \lambda_{\ell+1} | \beta \rangle. \quad (1.93)$$

A diagonalized unitary matrix is of the form,

$$\Gamma(C) = \begin{pmatrix} e^{i\gamma_1} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & e^{i\gamma_D} \end{pmatrix}. \quad (1.94)$$

It follows that,

$$\det \Gamma(C) = e^{i \sum_{\alpha=1}^D \lambda_\alpha} \quad (1.95)$$

$$= \det (\prod_{\ell=0}^{N-1} M_\ell). \quad (1.96)$$

The total Berry phase would be,

$$\sum_{\alpha=1}^D \lambda_\alpha = \text{Im} \ln \det (\prod_{\ell=0}^{N-1} M_\ell). \quad (1.97)$$

Exercise

1. Replace the single-valued snapshot states $|n, \lambda\rangle$ with $|n, \lambda'\rangle = e^{i\gamma_n(t)} |n, \lambda\rangle$, which are not necessarily single-valued. Show that

$$\langle n, \lambda' | \frac{\partial}{\partial t} | n, \lambda' \rangle = 0. \quad (1.98)$$

A state that evolves under such a restriction is said to be parallel transported. A state that moves under such **parallel transport condition** acquires a Berry phase $\gamma_n(C)$ after a cycle C .

2. For a spin-1/2 electron in a magnetic field $\mathbf{B} = B\hat{n}$, $\hat{n} = (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$, the eigenstates are,

$$|\hat{n}, +\rangle = \begin{pmatrix} \cos \frac{\theta}{2} \\ e^{i\phi} \sin \frac{\theta}{2} \end{pmatrix}, \quad |\hat{n}, -\rangle = \begin{pmatrix} -e^{-i\phi} \sin \frac{\theta}{2} \\ \cos \frac{\theta}{2} \end{pmatrix}.$$

(a) Show that the Berry connections are,

$$\mathbf{A}_\pm(\mathbf{B}) = \mp \frac{1}{2B} \frac{1 - \cos \theta}{\sin \theta} \hat{e}_\phi.$$

(b) Show that the Berry curvatures are,

$$\mathbf{F}_\pm(\mathbf{B}) = \mp \frac{1}{2} \frac{\hat{B}}{B^2}.$$

3. An electron moving in a magnetic field $\mathbf{B}(\mathbf{r}, t)$ has the Hamiltonian,

$$H = \frac{p^2}{2m} + \mathbf{B}(\mathbf{r}, t) \cdot \boldsymbol{\sigma}. \quad (1.99)$$

Suppose the eigenstates of the Zeeman term are $|n, \mathbf{B}(\mathbf{r})\rangle$,

$$\mathbf{B}(\mathbf{r}, t) \cdot \boldsymbol{\sigma} |n, \mathbf{B}(\mathbf{r}, t)\rangle = \varepsilon_n |n, \mathbf{B}(\mathbf{r}, t)\rangle, \quad (1.100)$$

expand the wave function of the moving electron as,

$$|\Psi\rangle = \sum_{n=\pm} \psi_n(\mathbf{r}, t) |n, \mathbf{B}\rangle. \quad (1.101)$$

From the Schrödinger equation, $H|\Psi\rangle = i\hbar \partial |\Psi\rangle / \partial t$, show that,

$$\left[\frac{1}{2m} (\mathbf{p} - \hbar \mathbf{A}_n)^2 + \hbar V_n + \varepsilon_n \right] \psi_n = i\hbar \frac{\partial \psi_n}{\partial t}, \quad (1.102)$$

in which

$$\mathbf{A}_n(\mathbf{r}, t) = +i \langle n, \mathbf{B} | \frac{\partial}{\partial \mathbf{r}} | n, \mathbf{B} \rangle, \quad (1.103)$$

$$V_n(\mathbf{r}, t) = -i \langle n, \mathbf{B} | \frac{\partial}{\partial t} | n, \mathbf{B} \rangle. \quad (1.104)$$

The off-diagonal coupling between $|+\rangle$ and $|-\rangle$ has been ignored (this is all right for a smooth magnetic field).

That is, the effective Hamiltonian for the particle motion acquires a Berry potential \mathbf{A}_n and a scalar potential V_n . Such potentials would result in forces (proportional to Berry curvatures) that act on the electron.

For example, the \mathbf{B} field here can be an effective magnetic field from a spin texture, such as a **magnetic skyrmion** (see [Nagaosa and Tokura, 2013](#)). An electron moving near a magnetic skyrmion would feel such forces.

4. For a spin-3/2 quadrupole coupling with a magnetic field, the projected Berry connections for the $|m| = 3/2$ sector and the $|m| = 1/2$ sector are,

$$\begin{aligned} \vec{A}^{(3/2)} &= \frac{1}{B \sin \theta} \begin{pmatrix} \frac{3}{2}(\cos \theta - 1) & 0 \\ 0 & -\frac{3}{2}(\cos \theta - 1) \end{pmatrix} \hat{e}_\phi, \\ \vec{A}^{(1/2)} &= \frac{1}{B} \begin{pmatrix} 0 & -ie^{-i\phi} \\ ie^{i\phi} & 0 \end{pmatrix} \hat{e}_\theta \\ &+ \frac{1}{B \sin \theta} \begin{pmatrix} \frac{1}{2}(\cos \theta - 1) & -\sin \theta e^{-i\phi} \\ -\sin \theta e^{i\phi} & -\frac{1}{2}(\cos \theta - 1) \end{pmatrix} \hat{e}_\phi. \end{aligned}$$

Show that the Berry curvatures are,

$$\mathbf{F}_{\theta\phi}^{(3/2)} = -\frac{3}{2} \frac{1}{B^2} \sigma_3,$$

$$\mathbf{F}_{\theta\phi}^{(1/2)} = +\frac{3}{2} \frac{1}{B^2} \sigma_3.$$

References

- Berry, M. V., 1984, Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences **392**(1802), 45.
- Bitter, T., and D. Dubbers, 1987, Phys. Rev. Lett. **59**, 251.
- Cheng, T.-P., and L.-F. Li, 1988, *Gauge Theory of elementary particle physics* (Oxford University Press).
- Li, T., L. Duca, M. Reitter, F. Grusdt, E. Demler, M. Endres, M. Schleier-Smith, I. Bloch, and U. Schneider, 2016, Science **352**(6289), 1094.
- Murakami, S., N. Nagaosa, and S.-C. Zhang, 2003, Science **301**(5638), 1348.
- Nagaosa, N., and Y. Tokura, 2013, Nature Nanotechnology **8**, 899.
- Schiff, L. I., 1972, *Quantum Mechanics* (McGraw-Hill, Inc), 3rd edition.
- Tycko, R., 1987, Phys. Rev. Lett. **58**, 2281.
- Wilczek, F., and A. Zee, 1984, Phys. Rev. Lett. **52**, 2111.
- Winkler, R., 2003, *Spin-orbit Coupling Effects in Two-Dimensional Electron and Hole Systems*, volume 191 (Springer).
- Zee, A., 1988, Phys. Rev. A **38**, 1.