



黑體輻射等同於空腔輻射，也等同於電磁波與一群物質一起達成的熱平衡狀態！






Cosmic Microwave Background Radiation CMB 宇宙背景輻射

Penzias and Wilson found CMB accidentally in 1965 and won the 1978 Nobel




Bell Lab Home of Transistor







The 15 meter Holmdel horn antenna at Bell Telephone Laboratories in Holmdel, New Jersey was built in 1959 for pioneering work in communication satellites for the NASA ECHO I.

Horn Antenna 觀測到來自宇宙、均勻而同向的微波Microwave。
稱為 **Cosmic Microwave Background Radiation CMB** 宇宙背景輻射。



電磁波以頻率或波長為特徵：


$$\lambda f = c$$



以微波為主的熱輻射非常冷！

CMB is a blackbody radiation at 2.725K。

大部分是微波，很冷。




完美的黑體輻射！

$$\lambda_{\max} \sim 10^{-3} \text{m}$$


$$\lambda_{\max} \cdot T = 2.898 \times 10^{-3} \text{ m} \cdot \text{K}$$

$$\lambda_{\max} \sim 10^{-3} \text{ m}$$


$$T \sim 3.0 \text{ K}$$



Cosmic Microwave Background Spectrum from COBE



CMB 在星空中所有的方向都是 3.0 K！稱為CMB 的同向性 Isotropy。




不同角度的微波來自彼此距離非常遙遠的地方。

在廣大的宇宙中，背景輻射如何維持同一溫度呢？


能彼此作用才能達成熱平衡，過去這些電磁波必須彼此很靠近！

沒想到科學家已經早有答案了。

天文學家在二十世紀初開始對星光作光譜分析。



Edwin Hubble 1889-1953



Hubble Telescope




Mount Wilson Observatory 1917






The 100-inch (2.5 m) **Hooker telescope** located at Mount Wilson Observatory, California, was the world's largest telescope from 1917 to 1949.



The mirror of the Hooker telescope on its way up the Mount Wilson Toll Road on a Mack Truck in 1917.




Workmen assembling the polar axis of the Hooker telescope.




Edwin Hubble Papers/Courtesy of Huntington Library, San Marino, Calif.


This glass side of a photographic plate shows where Hubble marked novae. The red VAR! in the upper right corner marks his discovery of the first Cepheid variable star — a star that told him the Andromeda galaxy isn't part of our Milky Way.




W. W. Campbell with a 36-inch spectroscope



Source: The University of Virginia





A two-prism spectrograph




Vesto Slipher & spectrograph

Hubble發現遠處星光的紅移現象redshift：星光的光譜頻率低於靜止原子的光譜頻率！




Vesto Slipher
1875-1969




3C 273 is a quasar located in the constellation Virgo. It was the first quasar ever to be identified. It is the optically brightest quasar in our sky and one of the closest with a redshift, z , of 0.158.

這是單一星體。第一個發現的Quasar。



星光的光譜頻率低於靜止原子光譜頻率！

Absorption lines in the visible spectrum of a supercluster of distant galaxies (right), as compared to absorption lines in the visible spectrum of the Sun (left). Arrows indicate redshift. Wavelength increases up towards the red and beyond (frequency decreases).




$$1 + z = \frac{\lambda_{\text{observed}}}{\lambda_{\text{emitted}}}$$

根據都卜勒效應： $z \sim \frac{v}{c}$ 速度遠小於光速時。

測量 z 就等於測量發光物體遠離的速度 v 。

z 相當方便，就用來表示星體離開我們的速度。

3C 273 is a quasar with a redshift, z , of 0.158. 遠離速度接近光速的五分之一。




早年看到的是星系Galaxy。

處女座

大熊座

Milton Humason
1891-197



長蛇座

根據電磁波的都卜勒效應，星體應該是離我們遠去！

遠方的銀河正離我們遠去，離開的速度可由紅位移的程度測得，速度各不相同！

這是上一頁的第一個光譜。

Messier 58 (also known as *M58* and *NGC 4579*) is an **intermediate barred spiral galaxy** with a weak inner ring structure located within the **constellation Virgo**, approximately 68 million **light-years** away from **Earth**.^{[9][10]} It was discovered by **Charles Messier** on **April 15, 1779** and is one of four barred spiral galaxies that appear in Messier's catalogue.^{[11][12][13][14][15][Note 1]} M58 is one of the brightest galaxies in the **Virgo Cluster**.^{[16][17]} From 1779 it was arguably (though unknown at that time) the farthest known astronomical object^[18] until the release of the **New General Catalogue** in the 1880s and even more so the publishing of **redshift values in the 1920s**.

Early observations [edit]

Charles Messier discovered Messier 58, along with the **elliptical galaxies Messier 59** and **Messier 60**, on April 15, 1779.^[14] M58 was reported on the chart of the **Comet of 1779** as it was almost on the same parallel as the star **Epsilon Virginis**.^{[11][19]} Messier described M58 as a very faint **nebula** in Virgo which would disappear in the slightest amount of light he used to illuminate the **micrometer wires**.^{[11][20]} This description was later contradicted by **John Herschel's** observations in 1833 where he described it as a very bright galaxy, especially towards the middle. Herschel's observations were also similar to the descriptions of both **John Dreyer** and **William Henry Smyth** who said that M58 was a bright galaxy, mottled, irregularly round and very much brighter toward the middle.^[11]



Messier 58



Observation data (J2000 epoch)

Constellation	Virgo ^[1]
Right ascension	12 ^h 37 ^m 43.522 ^s ^[2]
Declination	+11° 49′ 05.498″ ^[2]
Redshift	0.00506 ^{[2][3]}
Heliocentric radial velocity	1517 ± 1 km/s ^{[2][3]}
Distance	21 megaparsecs (68 million light-years) ^{[2][4]}
Apparent magnitude (V)	9.7 ^[5]

Messier 87

🌐 60 languages

Article [Talk](#)

[Read](#) [Edit](#) [View history](#) [Tools](#)

From Wikipedia, the free encyclopedia


Coordinates:  12°30′49.4″S, +12°23′28″E﻿ / ﻿−12.5136111°S 12.3911111°E﻿ / -12.5136111; 12.3911111 

Messier 87 (also known as **Virgo A** or **NGC 4486**, generally abbreviated to **M87**) is a **supergiant elliptical galaxy** in the **constellation Virgo** that contains several trillion stars. One of the **largest and most massive** galaxies in the local universe,^[b] it has a large population of **globular clusters**—about 15,000 compared with the 150–200 orbiting the **Milky Way**—and a jet of energetic **plasma** that originates at the core and extends at least 1,500 **parsecs** (4,900 **light-years**), traveling at a **relativistic speed**. It is one of the brightest radio sources in the sky and a popular target for both amateur and professional **astronomers**.

The French astronomer **Charles Messier** discovered M87 in 1781, and cataloged it as a **nebula**. M87 is about 16.4 million parsecs (53 million light-years) from Earth and is the second-brightest galaxy within the northern **Virgo Cluster**, having many **satellite galaxies**. Unlike a disk-shaped **spiral galaxy**, M87 has no distinctive **dust lanes**. Instead, it has an almost featureless, **ellipsoidal** shape typical of most **giant elliptical galaxies**, diminishing in **luminosity** with distance from the center. Forming around one-sixth of its mass, M87's **stars** have a nearly spherically symmetric distribution. Their population density decreases with increasing distance from the core. It has an active **supermassive black hole** at its core, which forms the primary component of an **active galactic nucleus**. The black hole was imaged using data collected in 2017 by the **Event Horizon Telescope** (EHT), with a final, processed image released on 10 April 2019.^[13] In March 2021, the EHT Collaboration presented, for the first time, a **polarized-based image** of the black hole which may help better reveal the forces giving rise to **quasars**.^[14]

The galaxy is a strong source of multi-wavelength radiation, particularly **radio waves**. It has an **isophotal** diameter of 40.55 kiloparsecs (132,000 light-years), with a diffuse galactic envelope that extends to a radius of about 150 kiloparsecs

Messier 87




Messier 87, with the blue **plasma jet** of its **galactic core** clearly visible (**composite image** of observations by the **Hubble Space Telescope** in **visible** and **infrared** light)

Observation data (J2000 epoch)

Constellation	Virgo
Right ascension	12 ^h 30 ^m 49.42338 ^s ^[1]
Declination	+12° 23′ 28.0439″ ^[1]
Redshift	0.00428 ± 0.00002 ^[2]
Heliocentric radial velocity	1,284 ± 5 km/s ^[2]
Distance	16.4 ± 0.5 Mpc (53.5 ± 1.6 Mly) ^[3]

這是上一頁列出的最遠的長蛇座星系。

LEDA 25177 (MCG+01-23-008)	1951-1960	$z=0.2$ ($V=61000$ km/s)	This galaxy lies in the Hydra Supercluster . It is located at B1950.0 $08^{\text{h}} 55^{\text{m}} 4^{\text{s}} +03^{\circ} 21'$ and is the BCG of the fainter Hydra Cluster Cl 0855+0321 (ACO 732). ^{[76][99][100][101][102][103][104][105]}
----------------------------	-----------	------------------------------	--



$$v \sim 0.2c$$

最大z值：14.32

JADES-GS-z14-0

🌐 5 languages

Article [Talk](#)

From Wikipedia, the free encyclopedia

JADES-GS-z14-0 is a [high-redshift Lyman-Break galaxy](#) in the [constellation Fornax](#) that was discovered in 2024 using [NIRCam](#) as part of the [JWST Advanced Deep Extragalactic Survey \(JADES\)](#) program.^{[1][2]} It has a redshift of 14.32, making it the most distant galaxy and astronomical object ever discovered.

Discovery [\[edit\]](#)

JADES-GS-z14-0 was observed using the James Webb Space Telescope's Near-Infrared Spectrograph (NIRSpec) in 2024,^[3] and it measured a redshift of 14.32,^[4] placing the galaxy's formation at an estimated 290 million years after the Big Bang.^[5] Its age, size, and luminosity added to a growing body of evidence that current theories of early star and galaxy formation are incomplete.^[6]

Characteristics [\[edit\]](#)

JADES-GS-z14-0 is 1600 light years wide and very luminous.^[6] Spectroscopic analysis revealed the presence of strong ionized gas emissions, including hydrogen and oxygen.^[4]

$$1 + z = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \left(1 + \frac{v}{c} \right)$$




JADES-GS-z14-0, as seen by NIRCam


Observation data (J2000 epoch)

Constellation	Fornax
Right ascension	03 ^h 32 ^m 36.89 ^s
Declination	−27° 46′ 49.33″
Redshift	14.32 ^{+0.08} _{−0.20}

Hubble發現銀河離開我們的速率，與該銀河與地球的距離成正比！1929




距離約百萬光年。



距離約千萬光年。

Galactic redshift vs. distance, plotted by Hubble and Humason (1931); red rectangle in lower left corner encloses data points plotted in 1929 graph above.




得到這個結果的努力中，距離的測量最難！



最近的數據，距離已經延伸至億光年。

Hubble發現銀河離開我們的速率，與該銀河與地球的距離成正比！

$$v = Hd \quad \text{Hubble's Law}$$



如果所有星系都以遠離地球的方向運動，而且越遠的星系，速度越快！
地球似乎應該是宇宙的中心！這太不可能了！

但理論天文物理學家對這個奇異的現象，其實已經有了答案。

Lemaître 勒梅特立刻認出這就是他的均勻宇宙膨脹的自然結果！



Lemaître et Eddington MIT 1932
autorisation de F. Laguens

Dear Professor Eddington

I just read the February No of the Observatory and the ~~discussion~~ on your suggestion of the investigating the ~~intermediate~~ non statistical intermediate solution between of Einstein's and de Sitter.

I made when investigation two years ago. I consider an universe of curvature constant in space but ~~variable~~ ^{increasing} with time. And I looked & emphasize the existence of a solution ~~for~~ ^{the} which the ~~apparent~~ ^{apparent} receding motion of the nebulae ~~is~~ ^{is} always ~~from~~ ^{from} a receding one ~~from~~ ^{from} time ~~to~~ ^{to} and from time minus infinity to ~~the~~ ^{the} infinity.

This solved the question put forward by de Sitter why the nebulae are on the receding branch of the hyperbola.


The result is as follows.

~~The~~ ^{The} receding motion of the nebulae is a measure of the initial or now assumed radius for $t = -\infty$. by the formulae

$$R_0 \approx \frac{2c}{\sqrt{3}} \quad \text{strictly} \quad \left[\frac{v}{2c} = \frac{1}{3R_0^2} + \frac{1}{R^2} - \frac{2}{3R_E^2} \right]$$

see later the work


Lemaître勒梅特神父是第一位提出擴張宇宙：Expanding Universe！



Georges Lemaître 1894-1966

Georges Lemaître


- 1896 - 1965
- Physicist and priest
- Father of cosmology
- He united Einstein's equations of general relativity and Hubble's observations
- Proposed the 'primeval atom theory' of the universe - a precursor to the 'The Big Bang' theory



Lemaître從愛因斯坦的解出發得出：宇宙的尺度是動態的！

宇宙的尺度是自然地隨時間而改變的。

讓我想像我們的宇宙，有一個個的星團。假設彼此相對大致是靜止的。
利用這些星團的位置為標記，可以在宇宙畫出尺格座標。
距離原則上可以測量，例如地球與下圖右的星團距離是 σ 。








重力根本不是力！

愛因斯坦提出：重力現象是質量造成周圍**時空彎曲**。

若無其他外力，所有物體都沿此**彎曲時空的直線**運動。

在彎曲時空中粒子走的直線由遠方看來是一條曲線！






彎曲時空代表時間、空間不是平坦的。



若平坦空間以棋盤狀等距尺格代表，彎曲空間以則扭曲的尺格來描述。

原來是平坦時空等距的格子，間距在彎曲時空下會被拉長或壓縮。




如果空間因重物而彎曲是可能的，……
空間像果凍一樣可以變形，那麼……

宇宙是靜止的呢？還是像果凍一樣、有彈性會來回搖晃……



彎曲空間中尺格可以隨空間位置而變化，但時間與空間是分不開的。
因此，尺格也可能會隨時間而變化(想像有彈力的彈簧的振盪)。

那麼兩個靜止的星系，彼此的距離，測量結果也可能跟時間有關！
換句話說，尺格可以是動態的。




尺格是動態的，**靜止星系**間的距離就會隨時間改變。

假設兩星系距離或尺格 σ 到了時間 t 時，變成原來時間 $t = 0$ 時距離 σ_0 的 $a(t)$ 倍。

$$\sigma(t) = a(t) \cdot \sigma_0$$

若宇宙是均勻的，所有地方的尺格變化 $a(t)$ 也會是均勻、即與位置無關。

所有的星系之間的距離，會以同樣的放大比例 $a(t)$ 來放大或縮小！



若所有的星系之間的距離，以同樣的放大比例 $a(t)$ 來放大！


$$a(t) > 1$$

那麼越遠的星系，自然遠離我們的速率越快！ $v = H \cdot d$

上圖左星系離我們的距離增加率，自然是上圖右星系距離增加率的兩倍。

簡言之，宇宙是均勻在擴張之中！

精細的計算：





若宇宙是均勻的，所有地方的尺度變化 $a(t)$ 也會是均勻而與位置無關。

$$\sigma(t) = a(t) \cdot \sigma(t = 0)$$

$$v = \frac{d}{dt} \sigma(t) = \frac{d}{dt} a(t) \cdot \sigma(0) = H \cdot a(t) \sigma(0) = H \cdot d$$


$$H = \frac{1}{a} \frac{da}{dt}$$

$$v = H \cdot d$$



哈伯定律可能顯示：宇宙正在擴張或膨脹之中，尺度放大比例在增加之中：

Expansion of the universe $a(t) \uparrow$




宇宙真得像果凍.....或麵包

MAP990404

$a(t) \uparrow$

我們怎麼知道放大比例 $a(t)$ 呢？

Alexander Friedmann 1888-1925



Friedmann 在1922 發現，尺度函數 $a(t)$ 可以由廣義相對論預測計算。
 $a(t)$ 滿足一個很簡單的微分方程式。

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = 8\pi GT_{\mu\nu}$$



$$\left(\frac{1}{a} \frac{da}{dt}\right)^2 = \frac{8\pi G}{3} \rho - \frac{K}{a^2}$$

Friedmann Equation

Cosmology的起點

尺度的變化率由當時宇宙的能量密度 ρ 、以及宇宙的曲度 K 決定！

簡言之，宇宙中的物質除了小範圍扭曲空間，造成重力效應。

也會大範圍的拉動空間尺格，使其隨時間變化。


比較奇妙的是，在廣義相對論下，空間還可以有一個自己的均勻彎曲度，
均勻的彎曲度也會影響尺度的變化率！

On the curvature of space


凸的

平的


凹的



Positively curved universe



Flat universe




Negatively curved universe

不同的彎曲度，它的效應可以以一曲度常數 K : $+1, 0, -1$ 表示。


$$\left(\frac{1}{a} \frac{da}{dt}\right)^2 = \frac{8\pi G}{3} \rho - \frac{K}{a^2}$$

現在暫時的測量結果是平的。




不同的彎曲度會造成宇宙不同的演化，擁有完全不同的未來！

Negative Curvature Models: $k = -1, \Omega_k > 0$ (infinite space)




Flat Models: $k = 0, \Omega_k = 0$ (infinite space)

$$\left(\frac{1}{a} \frac{da}{dt}\right)^2 = \frac{8\pi G}{3} \rho - \frac{K}{a^2} + \frac{\Lambda}{3}$$




Positive Curvature Models: $k = 1, \Omega_k < 0$, (finite space)




Classification of Friedmann Models

考慮物質含量及宇宙常數 Λ 的話，可能性就更多了。

重點是在所有可能性之中，幾乎沒有維持不變的的可能！ $a(t) \neq$ 常數





除非曲度常數正好為零，宇宙常數與能量密度正好抵消。





廣義相對論預測動態宇宙是非常自然的！所有星團距離都等比例變化。
放大比例 $a(t) = \text{常數}$ 的情況極為罕見，靜態宇宙是不自然且不穩定的！

這樣的想法是愛因斯坦無法接受的！



The results concerning the non-stationary world, contained in Friedmann's paper, appear to me suspicious 可疑. In fact, it turns out the solution given in it does not satisfy the equation of GR. 在這篇論文中的解釋不滿足廣義相對論的！


作為數學家的Friedmann 自然無法接受！立刻回信反駁。



愛因斯坦後來承認自己的錯誤: I am convinced that Mr. Friedmann's result are both correct and clarifying.

但愛因斯坦堅持他的偏見，在他的澄清信函的草稿中，還留有以下：
”a physical relevance can hardly be ascribed（很難看出與物理現實有甚麼
關聯）”這樣的字句。後來刪去了。


但因為愛因斯坦的意見，Friedmann的文章有十年完全被忽視。



當勒梅特在1927年告訴愛因斯坦他的動態宇宙的結果，

愛因斯坦回答：從物理的角度，這個理論在他看來十分abominable

（disgusting 令人厭惡的）他還無法忘記他對靜態宇宙的偏見。



Albert A. Michelson (center) with (left to right) M.L. Humason, Edwin Hubble, C.E. St. John, Albert Einstein, W.W. Campbell, and W.S. Adams in the library of the Mount Wilson Observatory, Pasadena, California, in early 1931.



On a visit to the Mount Wilson Observatory near Pasadena in 1931 :




愛因斯坦最後在證據之前，才接受宇宙是動態的事實！

宇宙膨脹最直接的可觀測現象就是光的紅外移。





但在廣義相對論中，這不是發光體或接收體運動造成的都卜勒效應。

畢竟星體相對於方格是靜止的，所以彼此的相對運動也是靜止的。




WRONG: Because receding galaxies are moving through space and exhibit a Doppler shift.

In the Doppler effect, a galaxy's movement away from the observer stretches the light waves, making them redder (*top*). The wavelength of light then stays the same during its journey through space (*middle*). The observer detects the light, measures its Doppler redshift and computes the galaxy velocity (*bottom*).



如果紅外移是星團移動造成，另一邊的星球會看到星團靠近，光會發生藍外移，在地球上往任何方向都從未看到藍外移。因此紅外移現象並不是來自星團移動。宇宙紅移是廣義相對論尺格加大時的自然結果！舞台持續改變，稱為重力紅外移。



空間座標間距會隨重力改變，時間座標的間距也會：

時間彎曲最特別的現象是時鐘的快慢與重力有關！

計算結果：高處的時鐘走得比低處的時鐘快。

地表距地心 r 處，時鐘測量的一段時間 Δt_r 小於無重力下時鐘測量的 Δt_∞ ：

$$\Delta t_r = \sqrt{1 - \frac{2GM}{c^2 r}} \Delta t_\infty = \frac{a_r}{a_\infty} \Delta t_\infty$$

From Tokyo Skytree's observatory, researchers prove time passes faster at a high altitude



A message reading "Together we can all win" is displayed on Tokyo Skytree after Prime Minister Shinzo Abe declared a state of emergency to fight the new coronavirus, in Tokyo on April 7. | REUTERS

KYODO

[PRINT](#) [SHARE](#)



[ARTICLE HISTORY](#) | APR 19, 2020

A Japanese team of researchers has shown that time at Tokyo Skytree's observatory — around 450 meters above sea level — passes four nanoseconds faster per day than at near ground level.

The finding, based on extremely precise "optical lattice clocks" that only go out of synch by one second every 16 billion years, proves Albert Einstein's general theory of relativity, which predicts that clocks in a strong gravitational

如何比較兩地的時鐘？利用從一處發出的光，傳到另一處時觀察。
光由低處傳到高處，光的週期(即一段時間 Δt_r)會變長，頻率會降低。
可見光會往較低頻的紅色移：**Gravitational Redshift 重力紅移**

$$1 + z \sim \frac{a_{\text{high}}}{a_{\text{low}}}$$




Gravitational Redshift

The final test proposed by Einstein in 1916 was the gravitational redshift. This was finally measured by the Pound-Rebka experiment in 1959 by firing gamma rays up and down a 22m tower at Harvard.

Measuring a frequency change of 1 part in 10^{15} , their measurement agreed with the Einstein prediction with an uncertainty of 10%.

Five years later, the accuracy was improved to a 1% agreement and now measurements can accurately agree to less than a percent accuracy.






- Performed in the tower at the University of Harvard
- Gamma rays of 14.4keV emitted from nuclear transition of iron-57
- Detected at base of 22.6m tower

Energy emitted $hf_{em} + mgH$
Energy detected hf_o

Harvard Tower Experiment



In just 22.6 meters, the fractional gravitational red shift given by

$$\nu = \nu_0 \left[1 + \frac{gh}{c^2} \right]$$

is just 4.92×10^{-15} , but the Mössbauer effect with the 14.4 keV gamma ray from iron-57 has a high enough resolution to detect that difference. In the early 60's physicists Pound, Rebka, and Snyder at the Jefferson Physical Laboratory at Harvard measured the shift to within 1% of the predicted shift.

By just using the expression for gravitational potential energy near the Earth, and using the m in the relativistic energy expression, the gain in energy for a photon which falls distance h is

$$\Delta E = mgh = \frac{E}{c^2} gh = \frac{14.4 \text{ keV}}{c^2} g \cdot 22.6 \text{ m}$$

$$\Delta E = 3.5 \times 10^{-11} \text{ eV}$$




哈伯定律觀察到的星光紅移是典型的重力紅移Gravitational Redshift !

時間尺格是隨空間尺格以同樣比例 a 放大 !

RIGHT: Because expanding space stretches all light waves as they propagate.


a_{then}



a_{now}





Galaxies hardly move through space, so they emit light with nearly the same wavelength in all directions [top]. The wavelength gets longer during the journey, because space is expanding. Thus, the light gradually reddens [middle and bottom]. The amount of redshift differs from what a Doppler shift would produce.



$$1 + z \sim \frac{a_{\text{now}}}{a_{\text{then}}}$$

光從圖中的星團發射後(想像前文的低處)，傳到地球被接收到(高處)的時候，時間尺格已經擴張了，測得時間變大。觀測的效應就如前文：頻率向低頻紅色偏。

RIGHT: Because expanding space stretches all light waves as they propagate.



LEDA 25177 (MCG+01-23-008)	1951–1960	z=0.2 (V=61000 km/s)
----------------------------	-----------	-------------------------

$$1 + z \sim \frac{a_{\text{now}}}{a_{\text{then}}}$$

z值也被用來代表星光發出的時候的尺格放大比例 a_{then} ，

z值越大， a_{then} 越小，越是久遠，z值甚至直接取代了宇宙歷史的紀年 t 。

JADES-GS-z14-0

最大z值：14.32

文A 5 languages ▾

Article [Talk](#)

[Read](#) [Edit](#) [View history](#) [Tools](#) ▾

From Wikipedia, the free encyclopedia

JADES-GS-z14-0 is a [high-redshift Lyman-Break galaxy](#) in the [constellation Fornax](#) that was discovered in 2024 using [NIRcam](#) as part of the [JWST Advanced Deep Extragalactic Survey \(JADES\) program](#).^{[1][2]} It has a redshift of 14.32, making it the most distant galaxy and astronomical object ever discovered.

Discovery [\[edit\]](#)

JADES-GS-z14-0 was observed using the James Webb Space Telescope's Near-Infrared Spectrograph (NIRSpec) in 2024,^[3] and it measured a redshift of 14.32,^[4] placing the galaxy's formation at an estimated 290 million years after the Big Bang.^[5] Its age, size, and luminosity added to a growing body of evidence that current theories of early star and galaxy formation are incomplete.^[6]

Characteristics [\[edit\]](#)

JADES-GS-z14-0 is 1600 light years wide and very luminous.^[6] Spectroscopic analysis revealed the presence of strong ionized gas emissions, including hydrogen and oxygen.^[4]




JADES-GS-z14-0, as seen by NIRCam

Observation data (J2000 epoch)

Constellation	Fornax
Right ascension	03 ^h 32 ^m 36.89 ^s
Declination	−27° 46′ 49.33″
Redshift	14.32 ^{+0.08} _{−0.20}


Big Bang 宇宙大霹靂

宇宙正在擴張之中，這事實確立之後：
往過去看，宇宙自然就是越來越限縮，
能量限縮在越來越小的範圍，因此一定是越來越熱！



Hot Big Bang 大霹靂

越是過去，宇宙越密，溫度越高



Georges Lemaître
1894-1966 Belgium



George Gamow
1904-1968 Russia, US




the Cosmic Egg exploding at the moment of the creation

WHAT KIND OF EXPLOSION WAS THE BIG BANG?


WRONG: The big bang was like a bomb going off at a certain location in previously empty space.

In this view, the universe came into existence when matter exploded out from some particular location. The pressure was highest at the center and lowest in the surrounding void; this pressure difference pushed material outward.



RIGHT: It was an explosion of space itself.

The space we inhabit is itself expanding. There was no center to this explosion; it happened everywhere. The density and pressure were the same everywhere, so there was no pressure difference to drive a conventional explosion.



大霹靂不是一個爆炸，因為沒有一個爆炸中心點。

大霹靂是一個均勻的宇宙擴張，因為早期宇宙能量密度極高，稱大霹靂。



86 Standard Cosmology



Fig. 3.11: The presently observable Universe at the Planck time, assuming $\hbar = 0.4$ (100 \times magnification).

早期宇宙只有這麼大！ $a(t)$ 非常小！


因為沒有外面，大小並不重要，但與現在比，極度擁擠，因此激烈作用！




當溫度極高，激烈熱運動的碰撞會將物質的組成成分撞擊分開，
 早期宇宙是一個如試管的、擁擠的、激烈的反應物！
 幸運的是對這一管反應物，我們非常了解它的成分與反應。
 如同在特定溫度下，我們了解理想氣體的速度分布。





原子核內的質子與中子，甚至其內的夸克也無法再束縛在一起。
和電子形成湯一樣密度很高的漿體。

Ylem primordial substance from which all matter is formed





Light and matter
are coupled






接近宇宙霹靂時，溫度很高的帶電的物質湯，會放出黑體輻射！

輻射與物質湯不斷碰撞交互作用，如空腔輻射與空腔的器壁，一直維持熱平衡。


所以在接近大霹靂時的宇宙中，輻射會呈現**黑體輻射**的樣式，

輻射的溫度就是物質湯的溫度。

因為物質湯是均勻的、意思是與位置無關，因此輻射也是**均勻而同向**。



Alpher, 1948



Dicke 1964

物理學家猜測，早期宇宙物質湯的黑體輻射會在現在留下可以觀測的痕跡。
宇宙背景輻射一觀測到，立刻被猜到這就是早期宇宙物質湯的黑體輻射的遺跡。

COSMIC BLACK-BODY RADIATION*

One of the basic problems of cosmology is the singularity characteristic of the familiar cosmological solutions of Einstein's field equations. Also puzzling is the presence of matter in excess over antimatter in the universe, for baryons and leptons are thought to be conserved. Thus, in the framework of conventional theory we cannot understand the origin of matter or of the universe. We can distinguish three main attempts to deal with these problems.

It has been pointed out by one of us (P. J. E. P.) that the observation of a temperature as low as 3.5°K , together with the estimated abundance of helium in the protogalaxy, provides some important evidence on possible cosmologies (Peebles 1965). This comes

Two of us (P. G. R. and D. T. W.) have constructed a radiometer and receiving horn capable of an absolute measure of thermal radiation at a wavelength of 3 cm. The choice

R. H. DICKE
P. J. E. PEEBLES
P. G. ROLL
D. T. WILKINSON

May 7, 1965
PALMER PHYSICAL LABORATORY
PRINCETON, NEW JERSEY

THE ASTROPHYSICAL JOURNAL

VOLUME 142

NOVEMBER 15, 1965

NUMBER 4

THE BLACK-BODY RADIATION CONTENT OF THE UNIVERSE AND THE FORMATION OF GALAXIES*

P. J. E. PEEBLES

Palmer Physical Laboratory, Princeton University, Princeton, N J.

Received March 8, 1965; revised June 1, 1965


ABSTRACT

A critical factor in the formation of galaxies may be the presence of a black-body radiation content of the Universe. An important property of this radiation is that it would serve to prevent the formation of gravitationally bound systems, whether galaxies or stars, until the Universe has expanded to a critical epoch. There is good reason to expect the presence of black-body radiation in an evolutionary cosmology, and it may be possible to observe such radiation directly.

The 2019 Physics Laureates



The 2019 Nobel Prize in Physics are awarded "for contributions to our understanding of the evolution of the universe and Earth's place in the cosmos", with one half to [James Peebles](#) "for theoretical discoveries in physical cosmology" and the other half jointly to [Michel Mayor](#) and [Didier Queloz](#) "for the discovery of an exoplanet orbiting a solar-type star."



Images tweeted by @NobelPrize

New perspectives on our place in the universe





The Nobel Prize in Physics 2019 rewards new understanding of the universe's structure and history, and the first discovery of a planet orbiting a solar-type star outside our solar system. This year's Laureates have contributed to answering fundamental questions about our existence. What happened in the early infancy of the universe and what happened next? Could there be other planets out there, orbiting other suns?

James Peebles took on the cosmos, with its billions of galaxies and galaxy clusters. His theoretical framework, which he developed over two decades, starting in the mid-1960s, is the foundation of our modern understanding of the universe's history, from the Big Bang to the present day. Peebles' discoveries have led to insights about our cosmic surroundings, in which known matter comprises just five per cent of all the matter and energy contained in the universe. The remaining 95 per cent is hidden from us. This is a mystery and a challenge to modern physics.





對於人在宇宙中的地位之嶄新觀點

背景輻射隨著宇宙的擴張漸漸冷卻，類似絕熱膨脹，到今天只剩 2.725K。
嚴格說這是一個等熵過程 $\Delta S = 0$ 。因為擴張是一個均勻同向的過程，。
絕熱膨脹後現在的背景輻射依舊維持接近大霹靂時的均勻同向及黑體的特性。




背景輻射的同向性可以以大霹靂理論解釋！
背景輻射成為大霹靂理論最重要的直接證據！

Penzias and Wilson 的測量因大氣吸收微波，十分不精確



Cobe (COsmic Background Explorer)

CMB大部分是微波，空氣可吸收，所以精密的測量必須在大氣層外進行





The Nobel Prize in Physics 2006
John C. Mather, George F. Smoot

Share this: [f](#) [g+](#) [t](#) [+](#) 9 [✉](#)

The Nobel Prize in Physics 2006



Photo: P. Izzo

John C. Mather

Prize share: 1/2



Photo: J. Bauer


George F. Smoot

Prize share: 1/2





The Nobel Prize in Physics 2006 was awarded jointly to John C. Mather and George F. Smoot *"for their discovery of the blackbody form and anisotropy of the cosmic microwave background radiation"*

Mather and George F. Smoot "for their discovery of the blackbody form and anisotropy of the cosmic microwave background radiation"

2.72548 ± 0.00057 K



宇宙背景輻射是均勻同向的完美的黑體，成為大霹靂理論最重要的直接證據！
也證實了從大霹靂到今天，近似是一個獨立的等熵冷卻的過程，不需要外力干預。




Mather and George F. Smoot "for their discovery of the blackbody form and **anisotropy** of the cosmic microwave background radiation"



Cobe 還有一個更重要的發現！

Cobe 同時觀察到大致同向的背景輻射有極微小的非同向性 anisotropy。


顏色代表冷熱，溫度差距大約是 $10^{-4} \sim 10^{-5} \text{K}$ 。

這個現象科學家也猜到了！





現在宇宙物質分布並不均勻，星團與星團之間是空的。
這應該起源於早期宇宙物質分布極微小的不均勻。
那背景輻射的溫度也應該會被影響而有微小的不均勻！



Peebles 1970



Zel'dovich 1970

LARGE-SCALE BACKGROUND TEMPERATURE AND MASS FLUCTUATIONS DUE TO SCALE-INVARIANT PRIMEVAL PERTURBATIONS

P. J. E. PEEBLES

Joseph Henry Laboratories, Physics Department, Princeton University

Received 1982 July 2; accepted 1982 August 13

ABSTRACT

The large-scale anisotropy of the microwave background and the large-scale fluctuations in the mass distribution are discussed under the assumptions that the universe is dominated by very massive, weakly interacting particles and that the primeval density fluctuations were adiabatic with the scale-invariant spectrum $P \propto \text{wavenumber}$. This model yields a characteristic mass comparable to that of a large galaxy independent of the particle mass, m_x , if $m_x \gtrsim 1$ keV. The expected background temperature fluctuations are well below present observational limits.

Subject headings: cosmic background radiation — cosmology — galaxies: formation


The expected temperature anisotropy at intermediate angular scales is given by equation (16). The rms fluctuation in T smoothed over $\theta = 10^\circ$ in a sample of size $\Theta = 100^\circ$ is

$$\delta T/T = w^{1/2} \sim 5 \times 10^{-6}. \quad (21)$$

由現在物質分布推算，背景輻射溫度不均勻度大約是 10^{-5} K。
正好與觀測結果相近。


宇宙的熱歷史

Thermal History of Universe



具體來說，背景輻射微小的不均勻是怎麼產生的呢？

為了回答這個問題，我們必須了解宇宙擴張的熱歷史！



宇宙是均勻的，所有地方的尺格放大比例 $a(t)$ 也是均勻而與位置無關。

$$\sigma(t) = a(t) \cdot \sigma(0)$$

時間尺格與空間尺格以同樣比例 a 放大！




Figure 5.8 The scale factor a as a function of time t (measured in units of the Hubble time), computed for the Benchmark Model. The dotted lines indicate the time of radiation–matter equality, $a_{rm} = 2.9 \times 10^{-4}$, the time of matter–lambda equality, $a_{m\Lambda} = 0.77$, and the present moment, $a_0 = 1$.

由Friedman Eq. 可以得到尺度放大比例與時間的關係： $a(t)$

宇宙的擴張是等熵過程。 $\Delta S = 0$

在早期的宇宙，以等熵膨脹計算，尺格放大比例與溫度成反比：

$$a \sim \frac{1}{T}$$


$$a(t) \sim \frac{1}{T} = 0$$

Pressure

而由Friedman Eq. 可以得到尺度大小與時間的關係： $a(t)$


背景輻射的溫度 T 與時間 t 對應，可以看成成宇宙歷史時間的指標！

$$t \uparrow \leftrightarrow T \downarrow$$




$$t \uparrow \leftrightarrow T \downarrow \leftrightarrow a \uparrow$$



Thermal History of Universe 而溫度決定了宇宙中物質的狀態！



快照



RIGHT: Because expanding space stretches all light waves as they propagate.



LEDA 25177 (MCG+01-23-008)	1951–1960	$z=0.2$ ($V=61000$ km/s)
----------------------------	-----------	------------------------------

$$1 + z \sim \frac{a_{\text{now}}}{a_{\text{then}}}$$

z 值也被用來代表星光發出的時候的尺格放大比例 a_{then} ，

z 值越大， a_{then} 越小，越是久遠， z 值甚至直接取代了宇宙歷史的紀年 t 。

JADES-GS-z14-0

最大z值：14.32

文A 5 languages ▾

Article [Talk](#)

[Read](#) [Edit](#) [View history](#) [Tools](#) ▾

From Wikipedia, the free encyclopedia


JADES-GS-z14-0 is a [high-redshift Lyman-Break galaxy](#) in the [constellation Fornax](#) that was discovered in 2024 using [NIRcam](#) as part of the [JWST Advanced Deep Extragalactic Survey \(JADES\)](#) program.^{[1][2]} It has a redshift of 14.32, making it the most distant galaxy and astronomical object ever discovered.

Discovery [\[edit\]](#)

JADES-GS-z14-0 was observed using the James Webb Space Telescope's Near-Infrared Spectrograph (NIRSpec) in 2024,^[3] and it measured a redshift of 14.32,^[4] placing the galaxy's formation at an estimated 290 million years after the Big Bang.^[5] Its age, size, and luminosity added to a growing body of evidence that current theories of early star and galaxy formation are incomplete.^[6]

Characteristics [\[edit\]](#)

JADES-GS-z14-0 is 1600 light years wide and very luminous.^[6] Spectroscopic analysis revealed the presence of strong ionized gas emissions, including hydrogen and oxygen.^[4]





JADES-GS-z14-0, as seen by NIRCam

Observation data (J2000 epoch)


Constellation	Fornax
Right ascension	03 ^h 32 ^m 36.89 ^s
Declination	−27° 46′ 49.33″
Redshift	14.32 ^{+0.08} _{−0.20}

Light and matter
are coupled



接近大霹靂時的黑體輻射與帶電的物質湯不斷碰撞作用，兩者一直維持熱平衡，
早期宇宙的能量以黑體輻射為主，物質猶如泡在熱湯 Heat Bath 內。

物質的溫度就是黑體輻射的溫度，也就是宇宙的溫度！



有了如此紀年方法，如果我們往回追溯，溫度 T 越來越高。 $t \downarrow \quad T \uparrow$


當溫度 T 足夠高時，熱運動的碰撞會將物質的組成成分撞擊分開。

例如溫度高於 $10^4\text{K} \sim 1\text{eV}$ ，電子會從原子中被撞擊與原子核分離。



溫度高於 $10^{10}\text{K} \sim 1\text{MeV}$ ，原子核中的質子和中子都無法束縛在一起。

因此每一段時間，宇宙物質的組成會不同，這就構成了宇宙的歷史！

Thermal History of the Universe





輻射能量與物質能量隨時間演化不一樣，後者會越來越重要！



早期宇宙輻射重於物質，是如在試管中、輻射液體內，擁擠激烈的反應物！
 幸運的是對這一管反應物，我們非常了解它的成分與反應的細節。
 如同在特定溫度下，我們了解理想氣體的速度分布。
 早期宇宙如同一個化學反應實驗！

早期宇宙是一團極濃的量子理想氣體！

雖然極濃，但因為溫度極高，粒子交互作用影響不大。



宇宙是處於熱平衡狀態！黑體輻射能量完全由溫度控制！

根據統計力學，各式粒子的數目也完全由溫度控制，可以計算出來。

早期宇宙是單面向的，非常簡單而容易預測！我們對它的成分一清二楚！

Bose-Einstein and Fermi-Dirac Distribution






Fig. 3.6: The equilibrium abundance of a species in a comoving volume element, $N = n/s$. Since both n_γ and s vary as T^3 , N is also proportional to n/n_γ .

粒子平衡時的數目，完全由溫度決定（此圖適用於黑體輻射以外的粒子）。

$$n_i(T) \sim g_i \left(\frac{m_i kT}{2\pi\hbar^2} \right)^{3/2} \exp\left(\frac{-m_i c^2 + \mu_i}{kT} \right)$$



現在順著歷史，時間增加，溫度下降，物質湯開始凝結。 $t \uparrow$ $T \downarrow$
 到了 10^{-6} s, 10^{13} K時，夸克開始組合成質子中子。熱能已不夠將其拆散。







一開始，溫度高時，由夸克生成的質子與中子的數量大致相等： $n_n \sim n_p$


溫度漸漸降低，夸克不再出現。中子質子不再由夸克生成。

質子與中子會透過與電子及微中子的反應互相轉換！



這個反應現在會繼續保持質子與中子處於熱平衡狀態！





$$\mu_e \sim \mu_\nu \sim 0$$

$$\mu_n \sim \mu_p$$

這個反應會保持質子與中子處於熱平衡狀態！

$$n_n \sim 2 \left(\frac{m_n kT}{2\pi\hbar^2} \right)^{3/2} \exp\left(\frac{-m_n c^2}{kT}\right)$$

$$n_p \sim 2 \left(\frac{m_p kT}{2\pi\hbar^2} \right)^{3/2} \exp\left(\frac{-m_p c^2}{kT}\right)$$

$$\frac{n_n}{n_p} \sim \exp\left[\frac{-(m_n - m_p)c^2}{kT}\right] = \exp\left(\frac{-Q}{kT}\right) \sim \exp\left(\frac{-1.29\text{MeV}}{kT}\right)$$

向右反應會放熱，

溫度下降，傾向向右反應： $n_n \downarrow$

直到 $t \sim 1\text{s}$ ， $kT \sim 0.8\text{MeV}$ ：

$$\tau(n + \nu \rightleftharpoons p + e^-) \ll \frac{1}{H}$$

空間擴張太快，此反應無法發生，
質子與中子數目凍結！

$$\frac{n_n}{n_p} \sim \exp\left(\frac{-1.29\text{MeV}}{0.8\text{MeV}}\right) \sim \frac{1}{5} = 0.2$$

Neutron Freeze Out or Neutrino Decoupling

根據比較精確的計算，這是發生在 $t \sim 20\text{s}$ ， $T \sim 1.3 \times 10^9\text{K}$ 。




Figure 9.2 Neutron-to-proton ratio in the early universe. The solid line assumes equilibrium; the dotted line gives the value after freezeout. Temperature decreases, and thus time increases, from left to right.





Figure 9.2 Neutron-to-proton ratio in the early universe. The solid line assumes equilibrium; the dotted line gives the value after freezeout. Temperature decreases, and thus time increases, from left to right.

根據精確的計算，在 $t \sim 20 \text{ s}$ ， $T \sim 1.3 \times 10^9 \text{ K}$ 時：

質子與中子數目凍結！注意中子少於質子！

$$\frac{n_n}{n_p} \sim \exp\left(\frac{-1.29 \text{ MeV}}{0.8 \text{ MeV}}\right) \sim \frac{1}{5} = 0.2$$


Neutron Freeze Out or Neutrino Decoupling




時間增加，溫度下降。 $t \uparrow$ $T \downarrow$

到了1s – 3 min, 10⁸ K時，質子中子開始組成氦原子核。這是極重要的時刻。

Big Bang Nucleosynthesis核合成



宇宙誕生三分鐘左右，熱能 $\sim 1 - 0.1$ MeV，質子與中子開始形成原子核。
核合成 Big Bang Nucleosynthesis BBN 就開始了。



質子與中子首先產生氫的同位素：氘 Deuterium的原子核。宇宙的首原子核！

Letters to the Editor

PUBLICATION of brief reports of important discoveries in physics may be secured by addressing them to this department. The closing date for this department is five weeks prior to the date of issue. No proof will be sent to the authors. The Board of Editors does not hold itself responsible for the opinions expressed by the correspondents. Communications should not exceed 600 words in length.

The Origin of Chemical Elements

R. A. ALPHER*

Applied Physics Laboratory, The Johns Hopkins University,
Silver Spring, Maryland

AND

H. BETHE

Cornell University, Ithaca, New York

AND

G. GAMOW

The George Washington University, Washington, D. C.

February 18, 1948

As pointed out by one of us,¹ various nuclear species must have originated not as the result of an equilibrium corresponding to a certain temperature and density, but rather as a consequence of a continuous building-up process arrested by a rapid expansion and cooling of the primordial matter. According to this picture, we must imagine the universe as a highly compressed state of general nuclear fluid and electrons which


We may remark at first that the building-up process was apparently completed when the temperature of the neutron gas was still rather high, since otherwise the observed abundances would have been strongly affected by the resonances in the region of the slow neutrons. According to Hughes,² the neutron capture cross sections of various elements (for neutron energies of about 1 Mev) increase exponentially with atomic number halfway up the periodic system, remaining approximately constant for heavier elements.

Using these cross sections, one finds by integrating Eqs. (1) as shown in Fig. 1 that the relative abundances of various nuclear species decrease rapidly for the lighter elements and remain approximately constant for the elements heavier than silver. In order to fit the calculated curve with the observed abundances³ it is necessary to assume the integral of $\rho_n dt$ during the building-up period is equal to 5×10^4 g sec./cm³.

On the other hand, according to the relativistic theory of the expanding universe⁴ the density dependence on time is given by $\rho \cong 10^6/t^2$. Since the integral of this expression diverges at $t=0$, it is necessary to assume that the building-up process began at a certain time t_0 , satisfying the relation:


$$\int_{t_0}^{\infty} (10^6/t^2) dt \cong 5 \times 10^4, \tag{2}$$

which gives us $t_0 \cong 20$ sec. and $\rho_0 \cong 2.5 \times 10^6$ g sec./cm³. This result may have two meanings: (a) for the higher densities existing prior to that time the temperature of the neutron gas that no... (b) of the uni... value... /cm³ which... if we




R. A. Alpher

α





H. Bethe

β




G. Gamow

γ



氕 Deuterium 氚 Tritium



這個生成反應如同一個化學反應！

宇宙猶如一個化學實驗的大熔爐！那就將化學反應的熱學直接移植過來即可！

$$\mu_p + \mu_n = \mu_D \quad \text{化學平衡的條件！}$$

由此條件可以得到氕的數目 n_D ，它是隨溫度而變化的，溫度越低， n_D 數目越多：

$$\frac{n_D}{n_p n_n} \propto T^{-\frac{3}{2}} \cdot e^{\frac{B_D}{T}}$$



$$\mu_p + \mu_n = \mu_D$$

化學平衡的條件！

$$n_D \sim g_D \left(\frac{m_D kT}{2\pi\hbar^2} \right)^{3/2} \exp\left(\frac{-m_D c^2 + \mu_D}{kT} \right)$$

此時這些粒子依舊處於熱平衡。

$$n_n \sim g_N \left(\frac{m_n kT}{2\pi\hbar^2} \right)^{3/2} \exp\left(\frac{-m_n c^2 + \mu_n}{kT} \right)$$

$$n_p \sim g_p \left(\frac{m_p kT}{2\pi\hbar^2} \right)^{3/2} \exp\left(\frac{-m_p c^2 + \mu_p}{kT} \right)$$

代入上三式計算 $\frac{n_D}{n_p n_n}$ ，根據化學平衡條件，化學能 μ_i 正好全部消去！

$$\frac{n_D}{n_p n_n} \propto T^{-\frac{3}{2}} \cdot e^{\frac{(m_p + m_n - m_D)c^2}{kT}} = T^{-\frac{3}{2}} \cdot e^{\frac{B_D}{kT}}$$


$$B_D \equiv (m_p + m_n - m_D)c^2 \sim 2.23 \text{ MeV}$$

氘的束縛能，也就是反應的放熱！

當 $T \gg B_D$ ， $n_D \sim 0$ 。當 $T \ll B_D$ ， $n_D \gg n_{p,n}$ 。


當熱能大於束縛能時，向左反應遠大於向右，束縛態氘就無法形成。

當熱能小於束縛能時，向左反應遠小於向右，束縛態氘會大量形成。



定溫定壓下的化學反應

定溫定壓的化學反應中，各氣體的分壓會隨隨反應進行而改變。



$$\Delta G = 0 \quad \text{平衡條件}$$

$$-\mu_{\text{N}_2} - 3\mu_{\text{H}_2} + 2\mu_{\text{NH}_3} = 0$$

$$-1 \left(\mu_{\text{N}_2,0} + RT \ln \frac{P_{\text{N}_2}}{P_0} \right) - 3 \left(\mu_{\text{H}_2,0} + RT \ln \frac{P_{\text{H}_2}}{P_0} \right) + 2 \left(\mu_{\text{NH}_3,0} + RT \ln \frac{P_{\text{NH}_3}}{P_0} \right) = 0$$

$$\ln \frac{P_{\text{H}_2}^3 P_{\text{N}_2}}{P_{\text{NH}_3}^2 P_0^2} = \frac{1}{RT} (2\mu_{\text{NH}_3,0} - \mu_{\text{N}_2,0} - 3\mu_{\text{H}_2,0}) \equiv \frac{\Delta G_0}{RT}$$

$$\frac{P_{\text{NH}_3}^2 P_0^2}{P_{\text{H}_2}^3 P_{\text{N}_2}} = e^{-\frac{\Delta G_0}{RT}}$$

這個條件決定了平衡時各個成分的分壓與分子數：

$$\frac{n_{\text{NH}_3}^2 n^2}{n_{\text{H}_2}^3 n_{\text{N}_2}} \cdot \frac{P_0^2}{P^2} = e^{-\frac{\Delta G_0}{RT}}$$



$$\frac{n_{\text{NH}_3}^2 n^2}{n_{\text{H}_2}^3 n_{\text{N}_2}} = e^{-\frac{\Delta G_0}{RT}}$$

$$\frac{n_D n_N}{n_p n_n} \propto T^{-\frac{3}{2}} \cdot e^{\frac{B_D}{T}}$$

ΔG_0 越大，溫度越低，越往右反應






Figure 9.3 The deuteron-to-neutron ratio during the epoch of deuterium synthesis. The


$$\frac{n_D}{n_p n_n} \propto T^{-\frac{3}{2}} \cdot e^{\frac{B_D}{T}}$$

大約 $t \sim 200 \text{ s}$ ， $T \sim 0.76 \times 10^9 \text{ K}$ ，就有相當足夠的氘生成了！






氦原子核 ${}^4\text{He}$



氦一產生後，會**非常快速地**再合成其他的原子核，最後會停頓在氦原子核 ${}^4\text{He}$ ：
 氦原子核 ${}^4\text{He}$ 比較穩定，很快地大部分的氘、氚及氦同位素 ${}^3\text{He}$ 都轉化為氦 ${}^4\text{He}$ ！
 氦 ${}^4\text{He}$ 再進一步製造更重的原子核就困難許多，量也很少！



10⁻⁴³ seconds
Temperature 10³²K
Gravity emerges

10⁻³⁵ seconds
Temperature 10²⁸K
Inflation era

10⁻⁴ seconds
Temperature 10¹³K
Antimatter disappears

THERMAL EQUILIBRIUM ERA

10⁻² seconds
10¹¹K

Equal numbers of protons and neutrons

1 billion photons for every proton or neutron

 Proton	 Neutron	 Antineutrino
 Electron	 Neutrino	 Positron



HYDROGEN ERA

1 second
10¹⁰K



HELIUM ERA

100 - 300 seconds
100,000 K

Almost 25% of visible universe is helium; 75% is hydrogen plus some tritium

Diagram illustrating the fusion chain for Helium-4 formation:

- Lithium-7 + photons → Helium-4 + Helium-4
- Helium-3 + Helium-4 → Beryllium-7 + photons
- Beryllium-7 + photons → Lithium-7
- Lithium-7 + Helium-4 → Beryllium-7 + photons
- Beryllium-7 + Helium-4 → Helium-3 + Helium-4
- Helium-3 + Helium-4 → Helium-4 + Helium-4

Other reactions shown include Tritium decaying into Helium-4 and Deuterium + photon forming Helium-4.


DEUTERIUM ERA

100 seconds
10⁹K



380,000 years
Atoms form.
Cosmic microwave background radiation permeates universe

100 - 200 million years
First stars form



^4He 會一直製造直到中子用盡為止！此時 Nucleosynthesis核合成就停止了。



^4He 產生很快，一般就以氦生成時間大約 $t \sim 200 \text{ s}$ 來標定核合成的時間！

因為氦原子核很穩定，絕大部分中子會進入 ^4He ，

我們可以由凍結的中子質子比，簡單估計可以形成的 ^4He 數量：

$$\frac{n_n}{n_p} \sim \frac{1}{5} \longrightarrow \frac{n_{^4\text{He}}}{n_N} \sim \frac{0.5}{6} \quad \frac{m_{^4\text{He}}}{m_N} \sim \frac{4 \cdot 0.5}{6} \sim \frac{1}{3} \quad \text{非常接近觀察結果：} \quad \frac{m_{^4\text{He}}}{m_N} \sim 0.25$$


考慮中子的衰變，此值的預測可以下修到：0.27。



真實的模型要複雜得多！但可以數值模擬計算！


中子耗盡後，核反應完全中斷，原子核不再與背景輻射達到熱平衡，

原子核種類數量固定直到現在。宇宙誕生三分鐘就決定了！



source: wikipedia

現在原子核種類數量對早期宇宙核子的絕對總數量非常敏感！







Big Bang Nucleosynthesis BBN 與觀察符合，成為大霹靂另一個有力證據

Essentially all of the elements that are heavier than lithium and beryllium were created much later, by stellar nucleosynthesis in evolving and exploding stars

BBN的預測成為大霹靂理論最重要的直接證據！

BBN能作如此簡單而精確的預測，是因為早期宇宙中，物質是處於熱平衡狀態！


這個預測，沒有被宇宙後來的演化破壞，也證實了從大霹靂到今天140億年間宇宙的擴張近似地是一個獨立的等熵的過程，不需要外力干預。



Recombination

宇宙中物質與宇宙背景輻射之間的熱平衡關係，並不是永久的。


Formation of light and matter



Frequent collisions between normal matter and light

Light and matter are coupled


Dark matter evolves independently: it starts clumping and forming a web of structures



As the Universe expands, particles collide less frequently

Light and matter separate


- Protons and electrons form atoms
- Light starts travelling freely: it will become the Cosmic Microwave Background (CMB)



Last scattering of light off electrons → **Polarisation**

Dark ages


Atoms start feeling the gravity of the cosmic web of dark matter



The Universe is dark as stars and galaxies are yet to form


First stars

The first stars and galaxies form in the densest knots of the cosmic web





Light from first stars and galaxies breaks atoms apart and "reionises" the Universe

Galaxy evolution



Light can interact again with electrons → **Polarisation**









Big Bang Nucleosynthesis BBN

Recombination
Photon Decoupling

大霹靂後約三十萬年，質子開始捕捉電子，形成原子
氫原子的束縛能大約~10eV

當熱能低於束縛能，氫原子就開始形成，而且無法拆散。

這又是一個宇宙歷史的里程碑。



大霹靂後約三十萬年，質子開始捕捉電子，形成原子，稱為**Recombination**。

$p + e^- \rightleftharpoons H + \gamma$ 捕捉的同時，拆散也還在進行，這是一個典型的化學平衡。

$$\frac{n_H}{n_p n_e} \propto T^{-\frac{3}{2}} \cdot e^{\frac{B_H}{kT}}$$

$$B_H \equiv m_p + m_e - m_H \sim 13.6 \text{ eV}$$

當熱能 kT 小於束縛能 B_H 時，向左反應遠小於向右，氫原子會大量形成。這時：

$$T_{\text{rec}} \sim 0.3 \text{ eV} \sim 3600 \text{ K}$$

$$t_{\text{rec}} \sim 288000 \text{ yr}$$




Figure 3.8: Free electron fraction as a function of redshift.


Recombination之後，電子數量急速降低↓！

$p + e^- \rightleftharpoons H + \gamma$ Recombination之後，向右反應遠快於向左！

電子密度急速降低！電中性的氫原子取而代之。

背景輻射與現在量較多的電中性的氫原子作用很小，

光子與電子的散射碰撞：



碰撞發生越來越趕不上宇宙擴張的速度： $\tau \gg \frac{1}{H}$


不久之後，光與物質便彼此獨立發展，稱為**Photon Decoupling**。

光在宇宙中幾乎自由行走，宇宙由模糊變透明。

從此，輻射的溫度不再是物質的溫度。這時：


$$T_D \sim 0.27\text{eV} \sim 3240\text{K}$$

$$t_D \sim 380000\text{yr}$$



Light and matter are coupled


Dark matter evolves independently: it starts clumping and forming a web of structures




As the Universe expands, particles collide less frequently

Light and matter separate

- Protons and electrons form atoms
- Light starts travelling freely: it will become the Cosmic Microwave Background (CMB)



Last scattering of light off electrons
→ **Polarisation**




Recombination is followed quickly by Photon Decoupling.

從此物質與背景輻射分道揚鑣，不再處於熱平衡。


物質與背景輻射會有各自的溫度。各自會有自己的故事。

Big Bang Nucleosynthesis BBN

Recombination
Photon Decoupling



宇宙此時密度如同空氣！



Light and matter separate

- Protons and electrons form atoms
- Light starts travelling freely; it will become the Cosmic Microwave Background (CMB)

Last scattering of light off electrons
→ **Polarisation**

Light and matter are coupled


Dark matter evolves independently; it starts clumping and forming a web of structures

As the Universe expands, particles collide less frequently

從此物質與背景輻射分道揚鑣，
 物質沒有非常均勻的背景輻射的干擾，才能發展出不均勻的分佈。
 若是沒有decoupling，現在整個宇宙還是一鍋均勻但稀疏的湯！

38萬年

Photon Decoupling




Photon Decoupling之前，背景輻射不斷被電子散射，宇宙是模糊的。
之後，背景輻射不再被散射而直進，宇宙由模糊變透明。
透明的宇宙對輻射不留下任何痕跡。

現在觀察到的背景輻射微波，自從宇宙誕生38萬年之後即無散射，而走直線。
它們的最後一次散射就發生在宇宙誕生38萬年時。

地球現在觀察到的微波在當時，分布於一個以地球為球心的球面上。

Surface of last scattering 最後散射面是位置的球面，也是歷史上特定的時間。




最後散射面對應的 z

在這球面上，這些光作了最後一次散射，


此球面上不同方向的物質分布的不均勻，就印在該角度的宇宙背景輻射微波上。

我們現在看到的宇宙背景輻射，猶如穿越雲層後被我們看到的光。
我們看到的光的影像，是雲層的下方表面，也就對應CMB的最後散射面！




背景輻射極微小的非同向性，就是這個球面上物質的不均勻分布。
CMB的非均勻性，保留了宇宙誕生38萬年時的影像！


Cobe 1991



宇宙誕生38萬年時，也就是138 億年前的影像！




宇宙早期微小的物質分布不均勻，演化到今日形成星球，以及星系分布的不均勻。





測量CMB非同向性，就是測量宇宙早期能量分布不均性。

宇宙背景輻射非同向性數據


WMAP 2005









PLANCK

Looking back to the dawn of time

Planck, ESA's time machine, is Europe's first mission that will look at the very edge of the observable Universe by studying the cosmic microwave background, the relic radiation of the Big Bang.

The radiation, which permeates space in all directions, is our deepest link to the birth of the Universe. It carries a picture of the universe as it was about 380 000 years after the Big Bang, or about 14 thousand million years ago, when light first made a small flight in space.

Thanks to our most advanced space experiment of its kind, the Planck telescope will measure tiny variations in the temperature of this otherwise unvarying background with the highest ever precision. These variations will reveal the fingerprints left by the seeds of the structures, such as galaxies, that will appear in our Universe today. With its sensitivity, Planck will reveal much more about the infant Universe than any mission has done so far.


Planck will help determine the properties of the Universe with unprecedented accuracy: the total density of matter and dark matter, the total amount of matter in the Universe, and the nature of dark energy.

The spacecraft carries a telescope and two panels of instruments designed to measure the microwave background. Their detectors are kept at perpetual zero close to liquid helium by a sophisticated cryogenic system.

Launched in 2009 on an Ariane 5ECA rocket from Europe's Spaceport in French Guiana, Planck will spend the next 5 years in orbit, located at about 1.5 million km from Earth, in the direction opposite of the Sun.

Life time is estimated at 15 months.








European Space Agency
 Agence spatiale européenne
 More information can be found on
<http://www.esa.int/planck>






The Planck one-year all-sky survey





(c) ESA, HFI and LFI consortia, July 2010

Planck 看到的星光




The anisotropies of the Cosmic Microwave Background (CMB) as observed by Planck 2013.

The Cosmic Microwave Background as seen by Planck and WMAP




WMAP



Planck

Planck 非常精密，對角度的解析更好



The Cosmos, Back in the Day
An image from data recorded by a European Space Agency satellite shows a heat map of the universe as it appeared 370,000 years after the Big Bang. Page A10.

Bronx Inspector, Secretly Taped, Suggests Race Is a Factor in Stops

By ANNEPASCALOTTEN
For years, the debate over the New York Police Department's use of stop-and-frisk tactics has centered on whether officers are guilty of racial profiling. Now, a recording suggests that, in at least one precinct, a person's skin color can be a deciding factor in who is stopped.

The recording, played on Thursday in Federal District Court in Manhattan, was of a conversation between a patrol officer and his commanding officer in the 147th Precinct in the South Bronx, a vicin-



Officer Pedro Sanchez

Fast-Growing Brokerage Firm Often Tangles With Regulators

Once Few, Women Hold More Power in Senat

By JENNIFER STEINBERGER
WASHINGTON — As her colleagues gathered for their first order of a new Congress, Senator Kelly Ayotte slipped into an empty Senate chamber to honor the grandeur of her legislative home. Ms. Ayotte, a Republican from New Hampshire, sat down at the wooden desk where generations of lawmakers from her state had used their desks, a deskman marched toward her with papers.


The desks, he stately told her, were for senators only.

Ms. Ayotte's induction that January day in 2011 into the most rarified ranks of the nation's political class — Senate senators — had begun. "The desk thing really stuck with me," Ms. Ayotte said. "There still just aren't that many of us."

In the 80 years since Rebecca Felton of Georgia became the first woman in the United States Senate — voters in her state in 1922 — women remain an anomaly in the upper chamber. But with 20 Senate senators now in office, an all-time high, women have risen from the curiosity they were for much of the 20th century into an important force



Universe as an Infant: Fatter Than Expected and Kind of Lumpy



European Space Agency; Planck Collaboration

A view of the cosmic microwave background collected by the European Space Agency's Planck satellite. The heat map of the cosmos was imprinted on the sky when the universe was just 380,000 years old.

By DENNIS OVERBYE
Published: March 21, 2013 | 210 Comments


Astronomers released the latest and most exquisite baby picture yet of the universe on Thursday, one that showed it to be 80 million to 100 million years older and a little fatter than previously thought, with more matter in it and perhaps ever so slightly lopsided.

Recorded by the European Space Agency's Planck satellite, the image is



Multimedia
Continued on Page A10




- FACEBOOK
- TWITTER
- GOOGLE+
- SAVE
- E-MAIL
- SHARE



Asymmetry in the average temperatures on opposite hemispheres of the sky





The 2015 *Planck* CMB temperature maps produced by all four methods (see an example in Fig. 7) are significantly more sensitive than those produced in 2013 (by a factor of 1.3).




The anisotropies of the Cosmic Microwave Background (CMB) as observed by Planck 2015.

我們會為了一張137億年的老照片這麼大費周章嗎？

這張照片除了當年的資訊，還包含了中間宇宙演化的歷史！







宇宙早期能量分布不均性，在擴張下演化為宇宙中的結構




Structure formation

宇宙早期物質密度微小的不均勻，由於引力吸引漸漸形成宇宙中的結構






星球才能形成





→ COSMIC HISTORY




我們對此宇宙早期微小的不均勻，有理論可以計算，等待驗證。



宇宙早期能量分布不均性，在最後散射面烙印於背景輻射之上，產生了CMB非同向性。

精確測量CMB非同向性即有機會驗證理論的計算。

而且CMB由當時傳播到現在、與物質雖無散射，但會受到中間宇宙擴張的影響。
CMB經歷了整個宇宙擴張的歷史。
CMB記錄了最後散射面上的不均勻度，同時包含了當時到現在宇宙演化的歷史，
兩者都對宇宙的特性十分敏感。



一般的做法就是設想一系列宇宙擴張的模型與參數，模擬預測出CMB的數據，再與觀測的數據對照，找出能正確預測數據的模型與參數！



例如非同向性的定量數據分析。

專業稱傅立葉分析，將左圖溫度差異對角度的變化作展開，得到右圖。

$$\Delta T(\phi) = \sum_l a_l \sin(l\phi)$$

意思就是，我們可以問左圖不均勻的顆粒粗細平均是多大。

$$\Delta T(\phi) = \sum_l a_l \sin(l\phi)$$

溫度差異對角度的變化作展開。變化度記為 l 。

較粗的點表示溫度相對角度變化較小，變化度 l 較小。








較細的點表示溫度相對角度變化較大，變化度 l 較大。

Sky Maps \rightarrow Power Spectra

We “see” the CMB sound as waves on the sky.

Shorter wavelengths are smaller frequencies are higher pitches

Use special methods to measure the strength of each wavelength.




Lineweaver 1997

這有點像畫畫：
將圖對色點的粗細做分解。




底色部分隨角度變化小，極粗。
倒影部分隨角度變化大，極細。



將CMB的非同向性分解為粗細的點的疊加：

紀錄不同粗細的點的比重：對角度的傅立葉分析出現明顯的峰值！

The Observed Sound Spectrum



PRIMEVAL ADIABATIC PERTURBATION IN AN EXPANDING UNIVERSE*

P. J. E. PEEBLES†

Joseph Henry Laboratories, Princeton University

AND

J. T. YU‡

Goddard Institute for Space Studies, NASA, New York

Received 1970 January 5; revised 1970 April 1

ABSTRACT

The general qualitative behavior of linear, first-order density perturbations in a Friedmann-Lemaître cosmological model with radiation and matter has been known for some time in the various limiting situations. An exact quantitative calculation which traces the entire history of the density fluctuations is lacking because the usual approximations of a very short photon mean free path before plasma recombination, and a very long mean free path after, are inadequate. We present here results of the direct integration of the collision equation of the photon distribution function, which enable us to treat in detail the complicated regime of plasma recombination. Starting from an assumed initial power spectrum well before recombination, we obtain a final spectrum of density perturbations after recombination. The calculations are carried out for several general-relativity models and one scalar-tensor model. One can identify two characteristic masses in the final power spectrum: one is the mass within the Hubble radius ct at recombination, and the other results from the linear dissipation of the perturbations prior to recombination. Conceivably the first of these numbers is associated with the great rich clusters of galaxies, the second with the large galaxies. We compute also the expected residual irregularity in the radiation from the primeval fireball. If we assume that (1) the rich clusters formed from an initially adiabatic perturbation and (2) the fireball radiation has not been seriously perturbed after the epoch of recombination of the primeval plasma, then with an angular resolution of 1 minute of arc the rms fluctuation in antenna temperature should be at least $\delta T/T = 0.00015$.






FIG. 4.—The residual mass-fluctuation spectrum $\mathcal{P}_m(k)^{1/2}$ (eq. [72]) in the open general-relativity model, $\rho_0 = 0.03\rho_c$ ($\rho_c = 1.8 \times 10^{-29} \text{ g cm}^{-3}$). The curve has been normalized to unity at maximum.



在 $l = 220, \theta \sim 1^\circ$ 左右有一個峯 peak

這表示CMB圖上冷熱之間最典型的角度是 1° 左右。

這對應最後散射面上能彼此聯絡、達成平衡、達成同溫的角度範圍。



在最後散射面上，宇宙只經歷了時間 t_{ls} ，
 在此面上能彼此有影響的區域最快是透過光，
 而到此時，光最多只能走 $\sim ct_{ls}$ ，這個範圍稱為 horizon $d_{Hor}(t_{ls})$ 。

$$\theta \sim \frac{d_{Hor}(t_{ls})}{d_A} \sim \frac{0.251\text{Mpc}}{12.8\text{Mpc}} \approx 0.020\text{rad} \approx 1.1^\circ$$

這區域演化傳播到現在地球，就對應到CMB圖上冷熱之間最典型的角 1° 左右。

峰的位置、高度，與宇宙的彎曲度，物質密度以及其他宇宙常數有關。
例如宇宙的彎曲度很明顯會影響地球上觀察CMB的視角。





Figure 5. The angular size of spots in the CMB are determined by the geometry.




① The first peak shows that the universe is geometrically flat, i.e. two parallel lines will never meet.

② The second peak shows that ordinary matter is just 5% of the matter and energy in the universe.


③ The third peak shows that 26% of the universe consists of dark matter.

From these three peaks, it is possible to conclude that if 31% (5%+26%) of the universe is composed of matter, then 69% must be dark energy in order to fulfil the requirement for a flat universe.


第一個峰值的位置正好是若宇宙為平的時後，理論計算預期的位置。
所以根據宇宙背景輻射的非同向性測量數據，宇宙是平的。




宇宙大尺度是平的！



大角度區 ΔT 是來自最後散射面上，暗物質分布不均勻所造成的重力位能脹落，進而影響CMB的光子。 ΔT 基本上是常數，被稱為平台，這是暴脹的特徵。



小角度區 ΔT 呈現一個個的peak，這是來自最後散射面上，在暗物質所產生的重力位能不均勻影響下，光子夸克電漿形成如音波般的駐波，進而影響CMB的光子。因為是駐波所以是一個一個的峰。



- ① The first peak shows that the universe is geometrically flat, i.e. two parallel lines will never meet.
- ② The second peak shows that ordinary matter is just 5% of the matter and energy in the universe.
- ③ The third peak shows that 26% of the universe consists of dark matter.

From these three peaks, it is possible to conclude that if 31% (5%+26%) of the universe is composed of matter, then 69% must be dark energy in order to fulfil the requirement for a flat universe.





Fig. 1. The *Planck* 2015 temperature power spectrum. At multipoles $\ell \geq 30$ we show the maximum likelihood frequency averaged temperature spectrum computed from the *Planck* cross-half-mission likelihood with foreground and other nuisance parameters determined from the MCMC analysis of the base Λ CDM cosmology. In the multipole range $2 \leq \ell \leq 29$, we plot the power spectrum estimates from the *Commander* component-separation algorithm computed over 94% of the sky. The best-fit base Λ CDM theoretical spectrum fitted to the *Planck* TT+lowP likelihood is plotted in the upper panel. Residuals with respect to this model are shown in the lower panel. The error bars show $\pm 1 \sigma$ uncertainties.

CMB的數據分析可以告訴我們宇宙的基本特性（參數）



95% confidence limits for the base Λ CDM model from *Planck* CMB power spectra, in combination with *Planck* lensing and external data (“ext,” BAO+JLA+ H_0). Nuisance parameters are not listed for brevity (they are in the *Planck* Archive tables), but the last three parameters give a summary measure of the total foreground or the three high- l temperature spectra used by the likelihood. In all cases the helium mass fraction prior mean $Y_p \approx 0.2453$, with theoretical uncertainties in the BBN predictions dominating over the

	TT+lowP+lensing 68% limits	TT+lowP+lensing+ext 68% limits	TT,TE,EE+lowP 68% limits	TT,TE,EE+lowP+lensing 68% limits	TT,TE,EE+lowP+lensing+ext 68% limits
23	0.02226 ± 0.00023	0.02227 ± 0.00020	0.02225 ± 0.00016	0.02226 ± 0.00016	0.02230 ± 0.00014
$\Omega_c h^2$	0.1197 ± 0.0022	0.1186 ± 0.0020	0.1184 ± 0.0012	0.1198 ± 0.0015	0.1188 ± 0.0010
$100\theta_{MC}$	1.04085 ± 0.00047	1.04103 ± 0.00046	1.04106 ± 0.00041	1.04077 ± 0.00032	1.04093 ± 0.00030
τ	0.078 ± 0.019	0.066 ± 0.016	0.067 ± 0.013	0.079 ± 0.017	0.066 ± 0.012
$\ln(10^{10} A_s)$	3.089 ± 0.036	3.062 ± 0.029	3.064 ± 0.024	3.094 ± 0.034	3.064 ± 0.023
n_s	0.9655 ± 0.0062	0.9677 ± 0.0060	0.9681 ± 0.0044	0.9645 ± 0.0049	0.9667 ± 0.0040
H_0	67.31 ± 0.96	67.81 ± 0.92	67.90 ± 0.55	67.27 ± 0.66	67.51 ± 0.64
Ω_Λ	0.685 ± 0.013	0.692 ± 0.012	0.6935 ± 0.0072	0.6844 ± 0.0091	0.6879 ± 0.0087
Ω_m	0.315 ± 0.013	0.308 ± 0.012	0.3065 ± 0.0072	0.3156 ± 0.0091	0.3121 ± 0.0087
$\Omega_m h^2$	0.1426 ± 0.0020	0.1415 ± 0.0019	0.1413 ± 0.0011	0.1427 ± 0.0014	0.1422 ± 0.0013
$\Omega_m h^3$	0.09597 ± 0.00045	0.09591 ± 0.00045	0.09593 ± 0.00045	0.09601 ± 0.00029	0.09596 ± 0.00030
σ_8	0.829 ± 0.014	0.8149 ± 0.0093	0.8154 ± 0.0090	0.831 ± 0.013	0.8150 ± 0.0087
$\sigma_8 \Omega_m^{0.5}$	0.466 ± 0.013	0.4521 ± 0.0088	0.4514 ± 0.0066	0.4668 ± 0.0098	0.4553 ± 0.0068
$\sigma_8 \Omega_m^{0.25}$	0.621 ± 0.013	0.6069 ± 0.0076	0.6066 ± 0.0070	0.623 ± 0.011	0.6091 ± 0.0067
z_{re}	$9.9^{+1.8}_{-1.6}$	$8.8^{+1.7}_{-1.4}$	$8.9^{+1.3}_{-1.2}$	$10.0^{+1.7}_{-1.5}$	$8.5^{+1.4}_{-1.2}$
$10^9 A_s$	$2.198^{+0.076}_{-0.085}$	2.139 ± 0.063	2.143 ± 0.051	2.207 ± 0.074	2.130 ± 0.053
$10^9 A_s e^{-2\tau}$	1.880 ± 0.014	1.874 ± 0.013	1.873 ± 0.011	1.882 ± 0.012	1.878 ± 0.011
Age/Gyr	13.813 ± 0.038	13.799 ± 0.038	13.796 ± 0.029	13.813 ± 0.026	13.807 ± 0.026
z_*	1090.09 ± 0.42	1089.94 ± 0.42	1089.90 ± 0.30	1090.06 ± 0.30	1090.00 ± 0.29
r_*	144.61 ± 0.49	144.89 ± 0.44	144.93 ± 0.30	144.57 ± 0.32	144.71 ± 0.31
$100\theta_*$	1.04105 ± 0.00046	1.04122 ± 0.00045	1.04126 ± 0.00041	1.04096 ± 0.00032	1.04106 ± 0.00031
z_{drag}	1059.57 ± 0.46	1059.57 ± 0.47	1059.60 ± 0.44	1059.65 ± 0.31	1059.62 ± 0.31
r_{drag}	147.33 ± 0.49	147.60 ± 0.43	147.63 ± 0.32	147.27 ± 0.31	147.41 ± 0.30
k_D	0.14050 ± 0.00052	0.14024 ± 0.00047	0.14022 ± 0.00042	0.14059 ± 0.00032	0.14044 ± 0.00032
z_{eq}	3393 ± 49	3365 ± 44	3361 ± 27	3395 ± 33	3382 ± 32
k_{eq}	0.01035 ± 0.00015	0.01027 ± 0.00014	0.010258 ± 0.000083	0.01036 ± 0.00010	0.010322 ± 0.000096
$100\theta_{s,eq}$	0.4502 ± 0.0047	0.4529 ± 0.0044	0.4533 ± 0.0026	0.4499 ± 0.0032	0.4512 ± 0.0031
f_{2000}^{143}	29.9 ± 2.9	30.4 ± 2.9	30.3 ± 2.8	29.5 ± 2.7	30.2 ± 2.7
$f_{2000}^{143 \times 217}$	32.4 ± 2.1	32.8 ± 2.1	32.7 ± 2.0	32.2 ± 1.9	32.8 ± 1.9
f_{2000}^{217}	106.0 ± 2.0	106.3 ± 2.0	106.2 ± 2.0	105.8 ± 1.9	106.2 ± 1.9



Table 9. Parameter 68 % confidence levels for the base Λ CDM cosmology computed from the *Planck* CMB power spectra, in combination with the CMB lensing likelihood (“lensing”).

Parameter	<i>Planck</i> TT+lowP+lensing
$\Omega_b h^2$	0.02226 ± 0.00023
$\Omega_c h^2$	0.1186 ± 0.0020
$100\theta_{MC}$	1.04103 ± 0.00046
τ	0.066 ± 0.016
$\ln(10^{10} A_s)$	3.062 ± 0.029
n_s	0.9677 ± 0.0060
H_0	67.8 ± 0.9 現在宇宙膨脹速度
Ω_m	0.308 ± 0.012 物質的佔比
$\Omega_m h^2$	0.1415 ± 0.0019
$\Omega_m h^3$	0.09591 ± 0.00045
σ_8	0.815 ± 0.009
$\sigma_8 \Omega_m^{0.5}$	0.4521 ± 0.0088
Age/Gyr	13.799 ± 0.038 現在宇宙的年紀
r_{drag}	147.60 ± 0.43
k_{eq}	0.01027 ± 0.00014 宇宙的曲度




Figure 21.1: Confidence level contours of 68.3%, 95.4% and 99.7% in the Ω_Λ - Ω_m plane from the Cosmic Microwave Background, Baryonic Acoustic Oscillations and the Union SNe Ia set, as well as their combination (assuming $w = -1$). Courtesy of Kowalski *et al.* [22]






Fig. 26. Constraints in the Ω_m - Ω_Λ plane from the *Planck* TT+lowP data (samples; colour-coded by the value of H_0) and *Planck* TT,TE,EE+lowP (solid contours). The geometric degeneracy between Ω_m and Ω_Λ is partially broken because of the effect of lensing on the temperature and polarization power spectra. These limits are improved significantly by the inclusion of the *Planck* lensing reconstruction (blue contours) and BAO (solid red contours). The red contours tightly constrain the geometry of our Universe to be nearly flat.



Before Planck



After Planck (2015)

背景輻射的圖像其實預測宇宙的總能量與質量中，有26%的黑暗物質，70%的黑暗能量，這兩個成分科學家都還完全不清楚它們究竟是什麼。所以我們所了解的部分，只佔宇宙總量的4%。所以有位教授是這樣說的：這場宇宙學的考試，科學家的得分只有四分，顯然是不及格的。但也顯示我們還處在科學發展的西部開墾時代，前方還有廣闊的知識土地，等待我們去探索。

Cosmic Confusion


Measurements of the universe don't agree on how fast it's expanding. Could an extra ingredient in the early cosmos explain the gap?

BY MARC KAMIONKOWSKI AND ADAM G. RIESS

ILLUSTRATION BY CHRIS GASH




**A Sampling of Hubble Constant Estimates,
Organized by Measuring Method**
Kilometers per second per 3.26 million light-years



RESULTS

Over the years the distance ladder measurements of the Hubble constant have converged at a value of 73 ± 1 kilometers per second per megaparsec (km/s/Mpc). The CMB method, on the other hand, gives an estimate of 67.5 ± 0.5 km/s/Mpc. The two values are too far apart to explain. Perhaps there is some overlooked error in the methods, or maybe they are telling us our cosmological model is incomplete.



The Dark Energy Spectroscopic Instrument (DESI) will spend five years creating a 3D map of the universe that will help reveal the nature of the dark energy driving cosmic expansion. The project's first six months of data show slivers of the universe that represent just 1 percent of the survey's ultimate volume of space. The colors represent different types of galaxies, including nearby bright galaxies in yellow, luminous red galaxies in magenta and galaxies with supermassive black holes in turquoise.

Credit: Source: "The Early Data Release of the Dark Energy Spectroscopic Instrument," by DESI Collaboration et al.; 2023 (data)

The Most Shocking Discovery in Astrophysics Is 25 Years Old

A quarter of a century after detecting dark energy, scientists are still trying to figure out what it is

By Richard Panek | December 1, 2023

One afternoon in early 1994 a couple of astronomers sitting in an air-conditioned computer room at an observatory headquarters in the coastal town of La Serena, Chile, got to talking. Nicholas Suntzeff, an associate astronomer at the Cerro Tololo Inter-American Observatory, and Brian Schmidt, who had recently completed his doctoral